

Law-Invariant Return and Star-Shaped Risk Measures*

Roger J. A. Laeven[†]

Dept. of Quantitative Economics
University of Amsterdam, CentER
and EURANDOM, The Netherlands
r.j.a.laeven@uva.nl

Emanuela Rosazza Gianin

Dept. of Statistics and Quantitative Methods
University of Milano-Bicocca, Italy
emanuela.rosazza1@unimib.it

Marco Zullino

Dept. of Mathematics and Applications
University of Milano-Bicocca, Italy
m.zullino@campus.unimib.it

This Version: May 9, 2024

Abstract

This paper presents novel characterization results for classes of law-invariant star-shaped functionals. We begin by establishing characterizations for positively homogeneous and star-shaped functionals that exhibit second- or convex-order stochastic dominance consistency. Building on these characterizations, we proceed to derive Kusuoka-type representations for these functionals, shedding light on their mathematical structure and intimate connections to Value-at-Risk and Expected Shortfall. Furthermore, we offer representations of general law-invariant star-shaped functionals as robustifications of Value-at-Risk. Notably, our results are versatile, accommodating settings that may, or may not, involve monotonicity and/or cash-additivity. All of these characterizations are developed within a general locally convex topological space of random variables, ensuring the broad applicability of our results in various financial, insurance and probabilistic contexts.

Keywords: Return risk measures; Star-shapedness; Law-invariance; SSD- and CSD-consistency; Value-at-Risk; Expected Shortfall.

MSC 2020 Classification: Primary: 91B06, 91B30, 60E15; Secondary: 60H30, 62P05.

JEL Classification: D81, G10, G20.

1 Introduction

Over the past decades, a large literature has developed the theory of monetary risk measures — monotone and cash-additive functionals — and analyzed their applications in a variety of fields including economics, finance, insurance, operations research and statistics. More recently, [5] introduced return risk measures — monotone and positively homogeneous functionals — and first results in the development of their static and dynamic theory and applications were obtained in [2, 5, 6, 30, 31]. Whereas monetary risk measures provide absolute assessments of risk, return risk measures provide relative assessments of risk, evocative of the distinct roles played by absolute and relative risk aversion measurements. The positive homogeneity property of return risk measures was relaxed by [30, 31] asserting the more general star-shapedness property.

*We are very grateful to the Editor and a referee for their comments and suggestions that have significantly improved this paper. This research was funded in part by the Netherlands Organization for Scientific Research under grant NWO Vici 2020–2027 (Laeven) and by an Ermenegildo Zegna Founder’s Scholarship (Zullino). Emanuela Rosazza Gianin and Marco Zullino are members of Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA, INdAM), Italy, and acknowledge the financial support of Gnampa Research Project 2024 (PRR-20231026-073916-203).

[†]Corresponding author.

Law-invariant risk measures — also referred to as law-determined or distribution-invariant risk measures — and stochastic order consistent risk measures play an important role in the theory and applications of risk measures.¹ This is due to their simplicity, tractability and statistical appeal, being statistical functionals. Many well-known risk measures are specific examples of law-invariant risk measures (e.g., Value-at-Risk, Expected Shortfall, the entropic risk measure and the p -norm). Law-invariant representations of coherent, convex, quasi-convex and quasi-logconvex risk measures have been derived in [10, 17, 25, 29, 30]. For stochastic order consistency of risk measures, we refer to, e.g., [3, 13, 35].

In this paper, we establish new characterization results for classes of law-invariant star-shaped functionals. The existing literature offers general representations for law-invariant convex and quasi-convex functionals that do not require monotonicity and/or cash-additivity; see the recent [4]. For a comprehensive discussion of non-monotone preferences and their applications, see, e.g., [1, 20, 41]. There is, however, a notable gap in the literature for star-shaped functionals, which we aim to fill. In [30], representations for law-invariant *quasi-logconvex* star-shaped risk measures are derived. Furthermore, in [9], representations for law-invariant *monetary* star-shaped risk measures are obtained, and in [28] the latter results are extended to allow for cash-subadditivity. By contrast, our representation results for law-invariant star-shaped functionals do not rely on the properties of monotonicity, quasi-logconvexity and/or cash-(sub)additivity. Furthermore, whereas [9, 28, 30] primarily focus on L^∞ as the space of random variables, we derive our results in a general locally convex topological space of random variables.

The generality and versatility of our setting — both in terms of properties that our functionals may satisfy and in terms of the spaces they are defined on — make some of the mathematical proofs intricate. Furthermore, while in the case of convex functionals the properties of law-invariance and convex-order stochastic dominance consistency are equivalent, provided that the functionals exhibit lower semicontinuity, such equivalence does not hold in general for star-shaped functionals, even not when assuming lower semicontinuity. Therefore, it becomes necessary to separately examine and analyze these two properties in the case of star-shaped functionals.

That is, the primary contribution of this paper is to bridge the gap between existing general representations for law-invariant convex and quasi-convex functionals on the one hand and those pertaining to star-shaped functionals on the other hand. We first establish representations for second-order and convex-order stochastic dominance consistent (SSD- and CSD-consistent, for short) star-shaped functionals. We also showcase the robustness of our proof strategies: when we impose the additional axioms of monotonicity and cash-additivity, our results align with those in [9], but within a (much) broader space than L^∞ . Second, we unveil novel representations à la Kusuoka [29] pertaining to SSD- and CSD-consistent star-shaped functionals, including their monetary variants that are previously unexplored in the literature. These characterization results may be viewed as a contribution of independent interest. Value-at-Risk and Expected Shortfall are pivotal building blocks in these representations. Third, we extend our analysis to general law-invariant star-shaped functionals as robustifications of Value-at-Risk. We demonstrate the validity of these representations even without the need for monotonicity or cash-additivity. Furthermore, in the case of law-invariant star-shaped risk measures that are also cash-additive, we present a more precise representation compared to that in Theorem 5 of [9], resulting in a reduced set over which minimization takes place. We demonstrate that these three sets of results extend to positively homogeneous functionals, thus generalizing [2, 30], and yielding in particular new representation results for SSD-consistent return risk measures. We finally provide three examples to illustrate that law-invariant star-shaped risk measures may arise naturally, also when cash-(sub)additivity is not preserved.

All the proof strategies share a common approach, relying on the approximation of star-shaped functions by convex ones. Consequently, our proof strategies effectively encapsulate the concept of star-shapedness, providing flexibility and adaptability in a myriad of settings

¹Law invariance is intimately related to probabilistic sophistication introduced by [34]; see also [36].

in which star-shapedness is assumed.

The remainder of this paper is organized as follows. In Section 2, we provide some preliminaries for law-invariant star-shaped functionals. In Section 3, we establish our first representation results for SSD- and CSD-consistent star-shaped functionals. Section 4 establishes representations à la Kusuoka [29] for SSD- and CSD-consistent star-shaped functionals. In Section 5, we derive representations as robustifications of Value-at-Risk for general law-invariant star-shaped functionals. Finally, Section 6 provides three illustrative examples.

2 Preliminaries

Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a non-atomic probability space and let L^0 be the space of real-valued random variables. Equalities and inequalities between random variables are meant to hold almost surely (a.s.). Given a set $\mathcal{X} \subseteq L^0$, we denote by \mathcal{X}_+ the subset of \mathcal{X} containing only positive random variables, i.e., $\mathbb{P}(X \geq 0) = 1$. The usual Lebesgue spaces of functions are denoted by L^p , $p \in [0, +\infty]$.

Definition 1. We write $X \sim Y$ if X and Y have the same law under \mathbb{P} . Furthermore, a set $\mathcal{X} \subseteq L^0$ is said to be law-invariant if for any $X, Y \in L^0$ with $X \in \mathcal{X}$ and $X \sim Y$ it holds that $Y \in \mathcal{X}$. When \mathcal{X} is a law-invariant set, a functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be law-invariant if, for any $X, Y \in \mathcal{X}$ with $X \sim Y$, we have $f(X) = f(Y)$.

We need the following assumptions on the spaces used in the sequel, which will be imposed throughout the paper unless explicitly mentioned otherwise. Let $\mathcal{X}, \mathcal{X}^*$ be two linear subspaces of L^0 and such that:

- $\mathcal{X}, \mathcal{X}^*$ are law-invariant;
- for any $X \in \mathcal{X}$ and $Y \in \mathcal{X}^*$ it holds that $XY \in L^1$;
- $L^\infty \subseteq \mathcal{X} \subseteq L^1$ and $L^\infty \subseteq \mathcal{X}^* \subseteq L^1$;
- \mathcal{X} contains the constants.

We consider the weakest topology $\sigma(\mathcal{X}, \mathcal{X}^*)$ on \mathcal{X} such that the linear functional $f_Y(X) := \mathbb{E}[XY]$ is continuous for every $Y \in \mathcal{X}^*$. Under these hypotheses, \mathcal{X} is a locally convex topological space. Additionally, for any $X \in \mathcal{X}$ and $c \in \mathbb{R}$, it holds that $X + c \in \mathcal{X}$, because \mathcal{X} is a linear space that includes constants. We will often use this fact without further mentioning. We recall that Orlicz spaces and L^p spaces with $p \in [1, +\infty]$ are examples of spaces satisfying the previous assumptions.

Definition 2. A functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ with $f(0) < +\infty$ is star-shaped if for any $\lambda \in (0, 1)$ and $X \in \mathcal{X}$ it holds that:

$$f(\lambda X) \leq \lambda f(X) + (1 - \lambda)f(0).$$

Furthermore, a functional $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ with $\rho(0) < +\infty$ is a risk measure if ρ is increasing w.r.t. the usual lattice order in L^0 , i.e., if for any $X, Y \in \mathcal{X}$ such that $X \geq Y$ it holds that $\rho(X) \geq \rho(Y)$. We define the proper domain of f as $\text{Dom}(f) := \{X \in \mathcal{X} : f(X) < +\infty\}$, and analogously for ρ .

From this point onward, we will assume that any functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ verifies $f(0) < +\infty$, i.e., $0 \in \text{Dom}(f)$, unless specified otherwise.

We note that, following the recent actuarial literature (see, e.g., [37]), Definition 2 adopts the sign convention that risk measures satisfy increasing rather than decreasing monotonicity. We recall that a functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is said to be lower semicontinuous if its level sets are $\sigma(\mathcal{X}, \mathcal{X}^*)$ -closed. In addition, if f is law-invariant and convex, and \mathcal{X} is

a rearrangement invariant space,² $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity is equivalent to the Fatou property (see Proposition 2.5 in [4]), i.e., for any sequence $(X_n)_{n \in \mathbb{N}} \subseteq \mathcal{X}$ and $X \in \mathcal{X}$ s.t. $X_n \rightarrow X$ a.s. with $\sup_{n \in \mathbb{N}} |X_n| \in \mathcal{X}$ it holds that $f(X) \leq \liminf_{n \rightarrow \infty} f(X_n)$. If f is monotone (thus, f is a risk measure) these two properties are also equivalent to continuity from below, i.e., for any $(X_n)_{n \in \mathbb{N}} \subseteq \mathcal{X}$ and $X \in \mathcal{X}$ such that $X_n \uparrow X$ a.s. we have $f(X) = \lim_{n \rightarrow \infty} f(X_n)$.

In the subsequent definition, we present a non-exhaustive set of axioms that a risk measure (or a functional) may satisfy.

Definition 3. Let $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a risk measure. Then ρ is said to be

- Normalized, if $\rho(0) = 0$.
- Convex, if $\rho(\lambda X + (1 - \lambda)Y) \leq \lambda \rho(X) + (1 - \lambda)\rho(Y)$, for any $\lambda \in (0, 1)$ and $X, Y \in \mathcal{X}$ with $X \neq Y$.
- Positively homogeneous, if $\rho(\lambda X) = \lambda \rho(X)$ for any $\lambda \in \mathbb{R}_+$ and $X \in \mathcal{X}$.
- Sublinear, if ρ satisfies convexity and positive homogeneity.
- Cash-additive, if $\rho(X + m) = \rho(X) + m$ for any $m \in \mathbb{R}$ and $X \in \mathcal{X}$.
- Cash-subadditive, if $\rho(X + m) \leq \rho(X) + m$ for any $m \in \mathbb{R}_+$ and $X \in \mathcal{X}$.

A monetary risk measure $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a risk measure that satisfies cash-additivity. Similarly, a return risk measure $\rho : \mathcal{X}_+ \rightarrow \mathbb{R} \cup \{+\infty\}$ is an increasing monotone functional verifying positive homogeneity for any $X \in \mathcal{X}_+$.³

The axioms in Definition 3 have been extensively studied in the literature (see, e.g., [14, 15, 18, 19, 21, 22, 24, 26, 32] and the references therein), also in connection with the law-invariance property (see, e.g., [4, 25, 29]).

Let us briefly revisit the core stochastic dominance concepts, particularly first-, second- and convex-order stochastic dominance. Under first-order stochastic dominance, we say that, for given $X, Y \in \mathcal{X}$, X dominates Y if $\mathbb{E}[g(X)] \geq \mathbb{E}[g(Y)]$ for every increasing function $g : \mathbb{R} \rightarrow \mathbb{R}$ such that the expectations exist; it is denoted as $X \succeq_1 Y$. Similarly, under second-order stochastic dominance, X dominates Y if $\mathbb{E}[g(X)] \geq \mathbb{E}[g(Y)]$ for every increasing and convex function $g : \mathbb{R} \rightarrow \mathbb{R}$; it is denoted as $X \succeq_2 Y$. Finally, X dominates Y in the convex order sense if $\mathbb{E}[g(X)] \geq \mathbb{E}[g(Y)]$ for every (not necessarily increasing) convex function $g : \mathbb{R} \rightarrow \mathbb{R}$; it is denoted as $X \succeq_c Y$. We use the symbol \sim to denote the relationship when both \succeq and \preceq apply simultaneously. Indeed, in this case, all relations \sim_1 , \sim_2 , and \sim_c coincide. Therefore, throughout the paper, we use the symbol \sim whenever any of these relations applies.

Definition 4. A risk measure $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is:

- FSD-consistent, if $\rho(X) \geq \rho(Y)$ for any $X, Y \in \mathcal{X}$ such that $X \succeq_1 Y$.
- SSD-consistent, if $\rho(X) \geq \rho(Y)$ for any $X, Y \in \mathcal{X}$ such that $X \succeq_2 Y$.

A functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is CSD-consistent if $f(X) \geq f(Y)$ for any $X, Y \in \mathcal{X}$ such that $X \succeq_c Y$.

Remark 5. We note that first-order stochastic dominance can be equivalently formulated as

$$X \succeq_1 Y \iff VaR_\beta(X) \geq VaR_\beta(Y) \quad \forall \beta \in [0, 1],$$

² \mathcal{X} is a rearrangement invariant space if it is a solid lattice with a law-invariant and complete lattice norm (for further details on rearrangement invariant spaces, cf. [12]).

³Canonical examples of return risk measures are Orlicz premia — Luxemburg norms from a purely mathematical perspective. Orlicz premia and their associated Orlicz spaces have been analyzed in a large literature in actuarial and financial mathematics (e.g., [5, 7, 8, 11, 16, 27, 33]).

where

$$\text{VaR}_\beta(X) := \inf\{x \in \mathbb{R} | \mathbb{P}(X \leq x) \geq \beta\}, \text{ for any } X \in \mathcal{X},$$

is the usual Value-at-Risk. Similarly, second-order stochastic dominance is equivalent to

$$X \succeq_2 Y \iff ES_\beta(X) \geq ES_\beta(Y) \quad \forall \beta \in [0, 1),$$

where, for any $X \in \mathcal{X}$,

$$ES_\beta(X) := \frac{1}{1-\beta} \int_\beta^1 \text{VaR}_m(X) dm, \text{ if } \beta \in [0, 1), \text{ and } ES_1(X) := \text{ess sup}(X),$$

is the usual Expected Shortfall.⁴ It is clear by this characterization that first-order stochastic dominance implies second-order stochastic dominance, i.e., if $X \succeq_1 Y$ then the relationship $X \succeq_2 Y$ also holds. Furthermore, convex-order stochastic dominance is equivalent to $X \succeq_2 Y$ with the further constraint $\mathbb{E}(X) = \mathbb{E}(Y)$, i.e., $ES_\beta(X) \geq ES_\beta(Y) \forall \beta \in [0, 1)$ and $\mathbb{E}(X) = \mathbb{E}(Y)$. Finally, we underline that if a functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is consistent with one of these three stochastic dominance conditions, then f is law-invariant. However, it is important to note that the converse implication does not hold, as we will explore in what follows. See [4, 22] for further details on this subject.

It is well-known that a functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ that adheres to either FSD- or SSD-consistency also displays monotonicity with respect to the standard pointwise order relation in L^0 . Therefore, FSD- or SSD-consistency are incompatible with functionals that lack monotonicity. In contrast to FSD- and SSD-consistency, the criterion of CSD-consistency does not entail monotonicity.

Remark 6. In this paper, we consistently present our findings for risk measures defined across the entire space \mathcal{X} . These findings encompass the setting of return risk measures defined on \mathcal{X}_+ as a specific case. This is justified because, given a positively homogeneous risk measure ρ on \mathcal{X}_+ , we can extend it to the whole space \mathcal{X} by defining:

$$\bar{\rho}(X) = \begin{cases} \rho(X) & \text{if } X \in \mathcal{X}_+, \\ +\infty & \text{otherwise,} \end{cases}$$

and

$$\rho'(X) := \inf\{\bar{\rho}(Y) : Y \in \mathcal{X}, Y \geq X\} = \inf\{\bar{\rho}(Y) : Y \in \mathcal{X}_+, Y \geq X\}, \quad (2.1)$$

with the convention $\inf\{\emptyset\} = +\infty$. It is clear that $\rho' : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is monotone and $\rho' = \rho$ on \mathcal{X}_+ . Moreover, it maintains positive homogeneity (and analogously for star-shapedness). Indeed, for any $\lambda \in (0, 1)$ and $X \in \mathcal{X}$, it holds that

$$\rho'(\lambda X) = \inf\{\bar{\rho}(Y) : Y \in \mathcal{X}_+, Y \geq \lambda X\} = \inf\{\bar{\rho}(\lambda Y') : Y' \in \mathcal{X}_+, Y' \geq X\} = \lambda \rho(X),$$

where we utilized positive homogeneity of ρ on \mathcal{X}_+ and the relation $\lambda Y \in \mathcal{X}_+ \iff Y \in \mathcal{X}_+$. Furthermore, if ρ is SSD-consistent, then ρ' inherits this property, as is evident by considering the relation $Y \succeq_2 X$ instead of $Y \geq X$ in Equation (2.1).

⁴It is worth noting that the relation $ES_\beta(X) \geq ES_\beta(Y)$ for all $\beta \in [0, 1)$ is equivalent to $ES_\beta(X) \geq ES_\beta(Y)$ for all $\beta \in [0, 1]$. Clearly, the second condition implies the first one. Moreover, for any $Z \in \mathcal{X} \subseteq L^1$, by Proposition 2.37 in [38] we have $ES_1(Z) = \lim_{\beta \rightarrow 1} ES_\beta(Z)$. Thus, it holds that: $ES_1(X) = \lim_{\beta \rightarrow 1} ES_\beta(X) \geq \lim_{\beta \rightarrow 1} ES_\beta(Y) = ES_1(Y)$. In the following, we will interchangeably use these two equivalent formulations of SSD.

3 SSD- and CSD-Consistent Star-Shaped Functionals

In this section, we explore the implications of the properties of SSD- and CSD-consistency for star-shaped risk measures within the framework of min-max representations. It has been shown in the literature (see e.g., [9]) that not all law-invariant star-shaped risk measures can be expressed as the minimum of law-invariant convex risk measures. This is exemplified by VaR, which falls into the category of star-shaped risk measures that cannot be represented as the minimum of law-invariant convex risk measures. The following theorem establishes the connection between SSD-consistent star-shaped risk measures and the minimum of SSD-consistent convex risk measures. Before proceeding further, we want to emphasize that the proof strategies employed in this section share a common thread with those provided in Section 4 of [31]. However, it is important to note that the setting here is fundamentally different. Unlike in [31], we do not rely on the order structure of \mathcal{X} or on the so-called C-property, which is pivotal in that context. Additionally, since our focus lies on CSD- and SSD-consistent functionals, we have to develop an alternative proof strategy to represent any star-shaped functional as the minimum of a convex family of functionals. In other words, it is not feasible to trivially adapt the proofs presented in [31] to achieve CSD- or SSD-consistent representations of star-shaped functionals.

Theorem 7. *A risk measure $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is SSD-consistent and star-shaped if and only if there exist a set of indices Γ and a family of SSD-consistent convex risk measures $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ with $\tilde{\rho}_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ such that $\tilde{\rho}_\gamma(0) = \rho(0)$ for all $\gamma \in \Gamma$ and*

$$\rho(X) = \min_{\gamma \in \Gamma} \tilde{\rho}_\gamma(X), \quad X \in \mathcal{X}. \quad (3.1)$$

In addition, $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a SSD-consistent and positively homogeneous risk measure if and only if there exists a family of SSD-consistent and sublinear risk measures $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ such that ρ can be represented as in Equation (3.1).

Proof. We provide a detailed proof for the star-shaped case, followed by a brief sketch of the proof for positively homogeneous risk measures, since it is similar to the star-shaped case.

The star-shaped case

‘Only if’: We start by proving that there exists a family of law-invariant convex functionals of which the pointwise minimum is ρ . Let us consider $\Gamma := \text{Dom}(\rho)$ and the family of functionals $(\rho_Z)_{Z \in \Gamma}$, whose elements $\rho_Z : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ are defined as:

$$\rho_Z(X) = \begin{cases} \alpha\rho(Z) + (1 - \alpha)\rho(0) & \text{if } \exists \alpha \in (0, 1] \text{ s.t. } X \sim \alpha Z, X \not\sim 0, \\ \rho(0) & \text{if } X \sim 0, \\ +\infty & \text{otherwise.} \end{cases}$$

ρ_Z is well-defined: If $X \sim 0$, this is trivially true. Note that the relation $X \sim c$ is equivalent to $X \equiv c$ for any $c \in \mathbb{R}$; we will leverage this observation without further mentioning. Let us assume $X \not\sim 0$. In this case, it is obvious that if $X \sim \alpha Z$ for some $\alpha \in (0, 1]$ then $Z \not\sim 0$. Now, we prove that such $\alpha \in (0, 1]$ is unique. To see this, let us assume that there exist $\alpha_1, \alpha_2 \in (0, 1]$ such that $X \sim \alpha_1 Z \sim \alpha_2 Z$, which holds if and only if:

$$\alpha_1 \text{VaR}_\beta(Z) = \alpha_2 \text{VaR}_\beta(Z) \quad \forall \beta \in [0, 1] \iff \alpha_1 = \alpha_2.$$

So, if $X \not\sim 0$, then $\alpha_1 = \alpha_2$. In particular, if $X \sim Y$, then either $X \sim Y \sim \alpha Z$ holds for some $\alpha \in (0, 1]$, yielding $\rho_Z(X) = \alpha\rho(Z) + (1 - \alpha)\rho(0) = \rho_Z(Y)$, or $X \sim Y$ but $X \not\sim \alpha Z$ for any $\alpha \in (0, 1]$, leading to $\rho_Z(X) = \rho_Z(Y) = +\infty$.

Properties of ρ_Z : We start by proving that $\rho_Z(X) \geq \rho(X)$ for any $X \in \mathcal{X}$. When $X \sim 0$ the assertion is trivial. We suppose $X \not\sim 0$. Let us assume there exists $\alpha \in (0, 1]$ such that

$X \sim \alpha Z$ (otherwise there is nothing to prove). Then, $\rho_Z(X) = \alpha\rho(Z) + (1 - \alpha)\rho(0) \geq \rho(\alpha Z) = \rho(X)$, by the SSD-consistency and star-shapedness of ρ . Furthermore, if $X \in \Gamma$, we can take $Z = X$, obtaining $\rho_X(X) = \rho(X)$. If $X \notin \Gamma$, then $\rho_Z(X) = +\infty$ for any $Z \in \Gamma$. Indeed, assuming, by contradiction, that $\rho_Z(X) < +\infty$, there exists $\alpha \in (0, 1]$ such that $\rho_Z(X) = \alpha\rho(Z) + (1 - \alpha)\rho(0) \geq \rho(\alpha Z) = \rho(X) = +\infty$, which leads to a contradiction. Thus, $\rho_Z(X) = \rho(X) = +\infty$. In summary, we have established that $\rho(X) = \min_{Z \in \Gamma} \rho_Z(X)$.

When $X \sim Y$, the law-invariance property of ρ_Z follows from the result established above. Indeed, if $X \sim Y \sim \alpha Z$ for some $\alpha \in (0, 1]$, then $\rho_Z(X) = \rho_Z(Y)$ by law-invariance of ρ . If $X \sim Y \not\sim \alpha Z$ for any $\alpha \in (0, 1]$, then $\rho_Z(X) = \rho_Z(Y) = +\infty$.

We are now prepared to define the family of SSD-consistent and convex risk measures $(\rho_Z)_{Z \in \Gamma}$. For any $Z \in \Gamma$, we set

$$\tilde{\rho}_Z(X) := \inf\{\rho_Z(Y) : Y \succeq_2 X\}, \quad X \in \mathcal{X}. \quad (3.2)$$

Properties of $\tilde{\rho}_Z$: $\tilde{\rho}_Z$ is SSD-consistent. Indeed, for any $X_1, X_2 \in \mathcal{X}$ with $X_1 \succeq_2 X_2$ we have that if $Y \succeq_2 X_1$ then $Y \succeq_2 X_2$, thus $\tilde{\rho}_Z(X_1) \geq \tilde{\rho}_Z(X_2)$. Hence, $\tilde{\rho}_Z$ is also monotone and law-invariant (see Remark 5). By definition of the infimum, it follows that $\tilde{\rho}_Z \leq \rho_Z$. Moreover, for each fixed $X \in \mathcal{X}$ and $Y \succeq_2 X$ it holds that $\rho_Z(Y) \geq \rho(Y) \geq \rho(X)$ (as shown in the first step of the proof), taking the infimum over $Y \succeq_2 X$ we obtain $\tilde{\rho}_Z(X) \geq \rho(X)$. Summing up, $\rho(X) \leq \tilde{\rho}_Z(X) \leq \rho_Z(X)$, hence $\rho(X) = \min_{Z \in \Gamma} \tilde{\rho}_Z(X)$. Now we prove convexity of $\tilde{\rho}_Z$. To see this, let us consider $X_1, X_2 \in \mathcal{X}$ with $X_1 \neq X_2$ and $\lambda \in (0, 1)$. If $\tilde{\rho}_Z(X_1) = +\infty$ or $\tilde{\rho}_Z(X_2) = +\infty$ there is nothing to prove. Moreover, if $\tilde{\rho}_Z(\lambda X_1 + (1 - \lambda)X_2) = +\infty$ then either $\tilde{\rho}_Z(X_1) = +\infty$ or $\tilde{\rho}_Z(X_2) = +\infty$. Indeed, if $\tilde{\rho}_Z(X_i) < +\infty$ for $i = 1, 2$, then there exist $\alpha_i \in [0, 1]$, $i = 1, 2$ such that $\tilde{\rho}_Z(X_i) = \alpha_i\rho(Z) + (1 - \alpha_i)\rho(0)$, $Y_i = \alpha_i Z \succeq_2 X_i$ and Y_i attains the infimum⁵ in the definition of $\tilde{\rho}_Z(X_i)$. This yields $(\lambda\alpha_1 + (1 - \lambda)\alpha_2)Z \succeq_2 \lambda X_1 + (1 - \lambda)X_2$, by positive homogeneity and subadditivity of Expected Shortfall. Thus by definition of $\tilde{\rho}_Z$ we have

$$\tilde{\rho}_Z(\lambda X_1 + (1 - \lambda)X_2) \leq (\lambda\alpha_1 + (1 - \lambda)\alpha_2)\rho(Z) + (1 - (\lambda\alpha_1 + (1 - \lambda)\alpha_2))\rho(0) < +\infty,$$

hence a contradiction. So, we only need to check the case $\tilde{\rho}_Z(\lambda X_1 + (1 - \lambda)X_2), \tilde{\rho}_Z(X_1), \tilde{\rho}_Z(X_2) < +\infty$. In this case we find at least one random variable $Y_i \succeq_2 X_i$ such that $Y_i \sim \alpha_i Z$, $\alpha_i \in [0, 1]$, $\rho_Z(Y_i) = \alpha_i\rho(Z) + (1 - \alpha_i)\rho(0)$ for $i = 1, 2$. Furthermore, it holds that

$$\begin{aligned} ES_\beta(\lambda X_1 + (1 - \lambda)X_2) &\leq \lambda ES_\beta(X_1) + (1 - \lambda)ES_\beta(X_2) \\ &\leq ES_\beta(\lambda Y_1) + ES_\beta((1 - \lambda)Y_2) = ES_\beta(\lambda\alpha_1 Z + (1 - \lambda)\alpha_2 Z), \end{aligned}$$

for any $\beta \in [0, 1]$. The first inequality follows from sublinearity of ES_β , the second inequality is due to the condition $Y_i \succeq_2 X_i$ for $i = 1, 2$, while the equality holds by positive homogeneity of ES_β , SSD-consistency of ES_β and the relation $Y_i \sim \alpha_i Z$ for $i = 1, 2$. Thus, $\lambda\alpha_1 Z + (1 - \lambda)\alpha_2 Z \succeq_2 \lambda X_1 + (1 - \lambda)X_2$ and

$$\begin{aligned} \tilde{\rho}_Z(\lambda X_1 + (1 - \lambda)X_2) &\leq \tilde{\rho}_Z(\lambda\alpha_1 Z + (1 - \lambda)\alpha_2 Z) \\ &\leq \rho_Z(\lambda\alpha_1 Z + (1 - \lambda)\alpha_2 Z) = \lambda\rho_Z(\alpha_1 Z) + (1 - \lambda)\rho_Z(\alpha_2 Z) \\ &= \lambda\rho_Z(Y_1) + (1 - \lambda)\rho_Z(Y_2), \end{aligned}$$

where the first inequality follows from SSD-consistency of $\tilde{\rho}_Z$, the second inequality follows from the relation $\tilde{\rho}_Z \leq \rho_Z$, the first equality is by definition of ρ_Z , while the definition of ρ_Z together with the relation $Y_i \sim \alpha_i Z$ with $i = 1, 2$ yields the second equality. Taking the infimum over $Y_i \succeq_2 X_i$ such that $Y_i \sim \alpha_i Z$ for some $\alpha_i \in [0, 1]$ and $i = 1, 2$ we get the thesis.

‘If’: We proceed to demonstrate the converse implication. Given a collection of SSD-consistent convex risk measures $(\rho_\gamma)_{\gamma \in \Gamma}$ that share the same value at 0 for all $\gamma \in \Gamma$, it is established that their pointwise minimum forms a star-shaped risk measure (refer to

⁵See Proposition 13 for a detailed proof of this statement.

Lemma 7 in [31]). Our focus now is to establish that $\rho(X) := \min_{\gamma \in \Gamma} \rho_\gamma(X)$ upholds SSD-consistency. To establish this, let us assume $X \succeq_2 Y$. By the SSD-consistency of the family $(\rho_\gamma)_{\gamma \in \Gamma}$, it follows that $\rho_\gamma(X) \geq \rho_\gamma(Y)$ for all $\gamma \in \Gamma$. This yields the result:

$$\rho(X) = \min_{\gamma \in \Gamma} \rho_\gamma(X) \geq \min_{\gamma \in \Gamma} \rho_\gamma(Y) = \rho(Y).$$

The positively homogeneous case

To prove the statement regarding positively homogeneous risk measures, we consider the family of functionals $(\rho_Z)_{Z \in \Gamma}$ defined as:

$$\rho_Z(X) = \begin{cases} \alpha \rho(Z) & \text{if } \exists \alpha \in (0, +\infty) \text{ s.t. } X \sim \alpha Z, X \not\sim 0 \\ 0 & \text{if } X \sim 0, \\ +\infty & \text{otherwise.} \end{cases}$$

By following the same reasoning as above, it can be proved that ρ_Z is well-defined for all $Z \in \Gamma$. In addition, $\rho_Z(X) \geq \rho(X)$ and $\rho_X(X) = \rho(X)$. Moreover, we observe that $\tilde{\rho}_Z := \inf\{\rho_Z(Y) : Y \succeq_2 X\}$ inherits the properties of ρ_Z and it is SSD-consistent and sub-linear. Thus, the family $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ with $\Gamma := \text{Dom}(\rho)$ fulfills the second thesis of the theorem. The converse implication is straightforward by observing that the pointwise minimum of sublinear risk measures is positively homogeneous. \square

Corollary 8. *$f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a CSD-consistent and star-shaped functional if and only if there exist a set of indices Γ and a family $(\tilde{f}_\gamma)_{\gamma \in \Gamma}$ of CSD-consistent and convex functionals $\tilde{f}_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ with $\tilde{f}_\gamma(0) = f(0)$ for all $\gamma \in \Gamma$, such that*

$$f(X) = \min_{\gamma \in \Gamma} \tilde{f}_\gamma(X), \quad X \in \mathcal{X}. \quad (3.3)$$

In addition, $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a CSD-consistent and positively homogeneous functional if and only if there exists a family of CSD-consistent and sublinear functionals $(\tilde{f}_\gamma)_{\gamma \in \Gamma}$ such that f can be represented as in Equation (3.3).

Proof. This can be proved similarly as in the proof of Theorem 7, recalling that the condition $\mathbb{E}(g(X)) \geq \mathbb{E}(g(Y))$ for any convex (not necessarily increasing) function $g : \mathbb{R} \rightarrow \mathbb{R}$ is equivalent to the relation $ES_\beta(X) \geq ES_\beta(Y)$ for all $\beta \in [0, 1]$ and $\mathbb{E}(X) = \mathbb{E}(Y)$, see Lemma 3 in [4]. Define:

$$f_Z(X) := \begin{cases} \alpha f(Z) + (1 - \alpha)f(0) & \text{if } \exists \alpha \in (0, 1] \text{ s.t. } X \sim \alpha Z, \\ f(0) & \text{if } X \sim 0, \\ +\infty & \text{otherwise,} \end{cases} \quad (3.4)$$

for any $Z \in \Gamma := \text{Dom}(f)$. Now, the proof follows verbatim from the proof of Theorem 7 considering $\tilde{f}_Z(X) := \inf\{f_Z(Y) : Y \succeq_c X\}$, $X \in \mathcal{X}$. \square

In addition to a comparison to the references discussed in the Introduction, the previous theorem and corollary might also be compared to the results obtained in Theorem 3.3 of [35]. There, the authors proved that a similar min-max representation holds in $\mathcal{X} = L^\infty$ when they consider SSD-consistent and *monetary* risk measures.

In the following proposition, we show that by requiring cash-additivity, we can obtain results akin to Theorem 4 in [9], but within the broader space \mathcal{X} . We acknowledge that it might be possible to derive the following results using continuity arguments and relying on Theorem 4 in [9]. However, we chose to pursue a different proof strategy, which shares the same underlying logic as that in Theorem 7, to demonstrate the robustness of this approach: it showcases how this strategy can encompass related results in the literature, while offering a distinct perspective.

Proposition 9. *A functional $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is an SSD-consistent, star-shaped and cash-additive risk measure if and only if there exist a set of indices Γ and a family $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ of SSD-consistent, convex and cash-additive risk measures $\tilde{\rho}_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ such that $\tilde{\rho}_\gamma(0) = \rho(0)$ for any $\gamma \in \Gamma$ and*

$$\rho(X) = \min_{\gamma \in \Gamma} \tilde{\rho}_\gamma(X), \quad X \in \mathcal{X}. \quad (3.5)$$

In addition, $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is an SSD-consistent, positively homogeneous and cash-additive risk measure if and only if there exist a set of indices Γ and a family $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ of SSD-consistent, sublinear and cash-additive risk measures $\tilde{\rho}_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ such that the representation in Equation (3.5) holds.

Proof. We establish the implication that if ρ is SSD-consistent, star-shaped and cash-additive, then there exists a family of SSD-consistent, convex and cash-additive risk measures whose pointwise minimum is ρ . The converse implication follows analogously to the proof of Theorem 7. We assume $\rho(0) = 0$ for brevity.

Let us define for any $Z \in \Gamma := \text{Dom}(\rho)$ the functional:

$$\rho_Z(X) := \begin{cases} \alpha\rho(Z) + c & \text{if } X \text{ is non-constant and there exist} \\ & \alpha \in (0, 1], c \in \mathbb{R} \text{ s.t. } X \sim \alpha Z + c, \\ c & \text{if there exists } c \in \mathbb{R} \text{ s.t. } X \sim c, \\ +\infty & \text{otherwise.} \end{cases}$$

Let us observe that if Z is constant, then the initial case in the definition of ρ_Z cannot occur. Indeed, when Z is constant, $ES_\beta(X) = ES_\beta(Z + c)$ holds for all $\beta \in [0, 1]$, and utilizing the cash-additivity of ES_β , we deduce $ES_\beta(X - Z - c) = 0$ for all $\beta \in [0, 1]$. By invoking the first part of the proof of Theorem 7, we conclude that $X = Z + c$ a.s., implying that X must be constant. Consequently, either X is constant and $\rho_Z(X) = c$, or $\rho_Z(X) = +\infty$. Moreover, ρ_Z is well-defined. Indeed, if Z is constant and there exists $c \in \mathbb{R}$ such that $X \sim c$, then $X = c$ a.s. and c is unique. If Z is not constant, suppose by contradiction there exist $\alpha_1, \alpha_2 \in [0, 1]$ and $c_1, c_2 \in \mathbb{R}$ such that $\alpha_1 \neq \alpha_2$ and $c_1 \neq c_2$. Then $ES_\beta(\alpha_1 Z + c_1) = ES_\beta(\alpha_2 Z + c_2)$, for any $\beta \in [0, 1]$. By cash-additivity and positive homogeneity of ES_β we have $ES_\beta(Z) = \frac{c_1 - c_2}{\alpha_1 - \alpha_2}$ for any $\beta \in [0, 1]$. Hence, the previous considerations yield $Z = \frac{c_1 - c_2}{\alpha_1 - \alpha_2}$ a.s., i.e., Z is constant, which is a contradiction. A similar conclusion holds if we suppose by contradiction $\alpha_1 = \alpha_2$ and $c_1 \neq c_2$ or $\alpha_1 \neq \alpha_2$ and $c_1 = c_2$. Furthermore, the cash-additivity of ρ_Z becomes evident. In particular, for any $m \in \mathbb{R}$, if $\rho_Z(X) < +\infty$, then $\rho_Z(X + m) = \alpha\rho(Z) + c + m = \rho_Z(X) + m$. This relation also holds when $\rho_Z(X) = +\infty$. Now, we proceed to demonstrate that $\rho_Z(X) \geq \rho(X)$ for all $X \in \mathcal{X}$, and $\rho_X(X) = \rho(X)$ for $X \in \Gamma$. Let us consider a non-constant $X \sim \alpha Z + c$ for some $\alpha \in (0, 1]$ and $c \in \mathbb{R}$ (the case where $X = c$ is clear due to the cash-additivity of ρ). In this setting, $\rho_Z(X) = \alpha\rho(Z) + c = \alpha\rho(Z + c/\alpha) \geq \rho(\alpha Z + c) = \rho(X)$, utilizing the cash-additivity and star-shapedness of ρ . Moreover, for $X \in \Gamma$, we can set $Z = X$, yielding $\rho_X(X) = \rho(X)$. Thus, for any $X \in \mathcal{X}$, we have $\rho(X) = \min_{Z \in \Gamma} \rho_Z(X)$. We observe that ρ_Z is law-invariant for any $Z \in \Gamma$. To see this, let us consider $X \sim X'$, thus $X \sim \alpha Z + c$ if and only if $X' \sim \alpha Z + c$, by law-invariance of ES_β , which implies $\rho_Z(X) = \rho_Z(X')$.

We can build a family of SSD-consistent, star-shaped and cash-additive risk measures $(\tilde{\rho}_Z)_{Z \in \Gamma}$ by introducing $\tilde{\rho}_Z(X) := \inf\{\rho_Z(Y) : Y \succeq_2 X\}$. In this case, the property of cash-additivity remains intact, as shown by

$$\tilde{\rho}_Z(X + m) = \inf\{\rho_Z(Y) : Y \succeq_2 X + m\} = \inf\{\rho_Z(W + m) : W \succeq_2 X\} = \tilde{\rho}_Z(X) + m,$$

where the last equality leverages the cash-additivity of ρ_Z and the definition of $\tilde{\rho}_Z$. Additionally, similarly to Theorem 7, we establish $\rho \leq \tilde{\rho}_Z \leq \rho_Z$, and also convexity of $\tilde{\rho}_Z$ can be checked similarly. As a result, the thesis is obtained. The thesis concerning positively

homogeneous risk measures is obtained by considering the family of sublinear functionals $(\rho_Z)_{Z \in \Gamma}$ defined by:

$$\rho_Z(X) := \begin{cases} \alpha \rho(Z) + c & \text{if } X \text{ is non-constant and there exist} \\ & \alpha \in (0, +\infty), c \in \mathbb{R} \text{ s.t. } X \sim \alpha Z + c, \\ c & \text{if there exists } c \in \mathbb{R} \text{ s.t. } X \sim c, \\ +\infty & \text{otherwise,} \end{cases}$$

and the corresponding family of SSD-consistent functionals $(\tilde{\rho}_Z)_{Z \in \Gamma}$ given by:

$$\tilde{\rho}_Z(X) := \inf\{\rho_Z(Y) : Y \succeq_2 X\}.$$

□

The following corollary extends our findings to CSD-consistent and star-shaped functionals, which are not necessarily monotone.

Corollary 10. *$f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a CSD-consistent, star-shaped and cash-additive functional if and only if there exist a set of indices Γ and a family $(f_\gamma)_{\gamma \in \Gamma}$ of CSD-consistent, convex and cash-additive functionals $f_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ with $f_\gamma(0) = f(0)$ for all $\gamma \in \Gamma$, such that*

$$f(X) = \min_{\gamma \in \Gamma} f_\gamma(X), \quad X \in \mathcal{X}. \quad (3.6)$$

In addition, $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a CSD-consistent, positively homogeneous and cash-additive functional if and only if there exist a set of indices Γ and a family $(f_\gamma)_{\gamma \in \Gamma}$ of CSD-consistent, sublinear and cash-additive functionals $f_\gamma : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ such that Equation (3.6) holds.

Proof. The proof follows verbatim from the proofs of Corollary 8 and Proposition 9. □

3.1 Representation results for Lipschitz risk measures

In the following results, we assume \mathcal{X} to be a rearrangement invariant space, since we need the structure of a normed space with a law-invariant norm. We will write $(\mathcal{X}, \|\cdot\|)$. We recall that a functional $f : \mathcal{X} \rightarrow \mathbb{R}$ is k -Lipschitz if there exists a $k > 0$ such that

$$|f(X) - f(Y)| \leq k\|X - Y\|, \quad \forall X, Y \in \mathcal{X}.$$

A family of functionals $(f_\gamma)_{\gamma \in \Gamma}$, where Γ is a set of indices, is said to be k -equi-Lipschitz if there exists a $k > 0$ such that:

$$|f_\gamma(X) - f_\gamma(Y)| \leq k\|X - Y\|, \quad \forall X, Y \in \mathcal{X} \text{ and } \gamma \in \Gamma.$$

Remark 11. *We note that a cash-subadditive risk measure can be derived from a k -Lipschitz risk measure by defining $\rho' = \rho/(k\|1\|)$. Conversely, any cash-subadditive risk measure is 1-Lipschitz as soon as the space \mathcal{X} has a unit. We refer to the proof of Lemma 5.1 in [28].⁶ This means that, on bounded random variables, cash-subadditivity is equivalent to a continuity condition. However, it is also worth noting that if ρ is defined on unbounded random variables, even in the case of law-invariant coherent risk measures, k -Lipschitz continuity may not necessarily arise from cash-(sub)additivity, as discussed in Remark 2.2 in [40].*

The next proposition gives a representation result for any SSD-consistent and star-shaped risk measure ρ that also verifies a k -Lipschitz condition. As a by-product, we obtain a representation result for SSD-consistent, star-shaped and cash-subadditive risk measures when \mathcal{X} has a unit.

⁶While the proof in [28] is presented for $\mathcal{X} = L^\infty$, it applies directly whenever \mathcal{X} has a unit, i.e., when \mathcal{X} consists of random variables with an order unit, equipped with the supremum norm.

Proposition 12. *Let $(\mathcal{X}, \|\cdot\|)$ be a rearrangement invariant space. Then $\rho : \mathcal{X} \rightarrow \mathbb{R}$ is a k -Lipschitz, SSD-consistent and star-shaped risk measure if and only if it can be represented as the pointwise minimum of a family $(\rho_\gamma)_{\gamma \in \Gamma}$ of k -Lipschitz, SSD-consistent and convex risk measures.*

In addition, if \mathcal{X} has a unit, $\rho : \mathcal{X} \rightarrow \mathbb{R}$ is an SSD-consistent, star-shaped and cash-subadditive risk measure if and only if it is the pointwise minimum of a family of convex, SSD-consistent and cash-subadditive risk measures.

Proof. The ‘if’ implication is trivial, as it is well-known that the pointwise minimum of k -equi-Lipschitz functionals remains k -Lipschitz.

We now demonstrate the ‘only if’ part. Let us fix $Z \in \mathcal{X}$ and consider $b_k(X) := k\|X\|$. We define

$$\begin{aligned} \rho^Z(X) &:= \inf_{\bar{X} \in \mathcal{X}} \{b_k(\bar{X}) + \tilde{\rho}_Z(X - \bar{X})\} \\ &= \inf_{\bar{X} \in \mathcal{X}} \{b_k(X - \bar{X}) + \tilde{\rho}_Z(\bar{X})\} \\ &= \inf \{k\|X - \bar{X}\| + \bar{\alpha}\rho(Z) + (1 - \bar{\alpha})\rho(0) : A_{\bar{X}} \neq \emptyset \text{ with } \bar{\alpha} = \min A_{\bar{X}}\}, \end{aligned}$$

where $\tilde{\rho}_Z$ is the SSD-consistent risk measure defined in Equation (3.2) and A_X ⁷ is defined in Equation (4.2). By definition, ρ^Z is the infimal convolution between $\tilde{\rho}_Z$ (convex and proper) and the Lipschitz and convex function $b_k(X)$. In particular, ρ^Z is the Pasch-Hausdorff envelope of $\tilde{\rho}_Z$, thus it is a proper, convex, and Lipschitz function. We claim that ρ^Z is monotone. Indeed, for any $X, Y \in \mathcal{X}$ with $X \geq Y$ it holds that $X - \bar{X} \geq Y - \bar{X}$ for all $\bar{X} \in \mathcal{X}$. Thus, monotonicity of $\tilde{\rho}_Z$ yields $b_k(\bar{X}) + \tilde{\rho}_Z(X - \bar{X}) \geq b_k(\bar{X}) + \tilde{\rho}_Z(Y - \bar{X})$ for any $\bar{X} \in \mathcal{X}$. Taking the infimum over $\bar{X} \in \mathcal{X}$ we get $\rho^Z(X) \geq \rho^Z(Y)$. We now prove that $\rho(X) = \min_{Z \in \mathcal{X}} \rho^Z(X)$ for any $X \in \mathcal{X}$. Fixing $X, \bar{X} \in \mathcal{X}$, with $A_{\bar{X}} \neq \emptyset$, we have:

$$\rho(X) - \bar{\alpha}\rho(Z) - (1 - \bar{\alpha})\rho(0) \leq \rho(X) - \rho(\bar{\alpha}Z) = \rho(X) - \rho(\bar{Y}) \leq \rho(X) - \rho(\bar{X}) \leq k\|X - \bar{X}\|.$$

Here, the first inequality follows from the star-shapedness of ρ , the equality stems from the law-invariance of ρ and the relation $\bar{Y} \sim \bar{\alpha}Z$ by the definition of $A_{\bar{X}}$, the second inequality is implied by the SSD-consistency of ρ and $\bar{Y} \succeq_2 \bar{X}$ by the definition of $A_{\bar{X}}$, while the last inequality holds by the k -Lipschitzianity of ρ . Rearranging and taking the infimum over the set $\{\bar{X} \in \mathcal{X} : A_{\bar{X}} \neq \emptyset\}$, we obtain $\rho^Z \geq \rho$. Moreover, $\rho^X(X) \leq \rho(X)$, by setting $X = \bar{X} = Z$ (which implies $\bar{\alpha} = 1$). Thus, the thesis follows.

It only remains to establish the SSD-consistency of ρ^Z . As ρ^Z is monotone, we can equivalently prove that ρ^Z is CSD-consistent. For this, it is sufficient to show the dilatation monotonicity property for ρ^Z . Indeed, as per Theorem 3.A.4 in [39], CSD-consistency holds if and only if we can prove that $\rho^Z(X) \geq \rho^Z(Y)$ for any $X, Y \in \mathcal{X}$ such that $Y = \mathbb{E}[X|Y]$. Given the assumption of a rearrangement-invariant space, b_k , being a norm multiplied by a scalar, is law-invariant, convex, and $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuous. Consequently, b_k satisfies CSD-consistency. Analogously, $\tilde{\rho}_Z$ is CSD-consistent, by Theorem 7. Now, let us consider $X, Y \in \mathcal{X}$ such that $Y = \mathbb{E}[X|Y]$. We have:

$$\begin{aligned} \rho^Z(X) &= \inf \left\{ b_k(X_1) + \tilde{\rho}_Z(X_2) : \sum_{i=1}^2 X_i = X \right\} \\ &\geq \inf \left\{ b_k(Y_1) + \tilde{\rho}_Z(Y_2) : \sum_{i=1}^2 Y_i = Y \right\} = \rho^Z(Y), \end{aligned}$$

where the inequality follows from CSD-consistency of b_k and $\tilde{\rho}_Z$, with $Y_i = \mathbb{E}[X_i|Y]$ for $i = 1, 2$, and it is evident that $Y_1 + Y_2 = Y$ since $X_1 + X_2 = X$. This shows the first thesis.

⁷In the proof of Proposition 13 it is demonstrated that if $A_X := \{\alpha \in [0, 1] : \exists Y \in \mathcal{X} : Y \succeq_2 X, Y \sim \alpha Z\} \neq \emptyset$ then there exists a (unique) minimizer $\bar{\alpha} \in A_X$ such that $\tilde{\rho}_Z(X) = \bar{\alpha}\rho(Z) + (1 - \bar{\alpha})\rho(0)$. Clearly, this also implies the existence of $\bar{Y} \in \mathcal{X}$ such that $\bar{Y} \succeq_2 X$ and $\bar{Y} \sim \bar{\alpha}Z$.

If \mathcal{X} has a unit, then ρ is cash-subadditive if and only if ρ is 1-Lipschitz continuous, by Remark 11. Consequently, the latter part of this proposition directly follows from the former. \square

In [28], it is demonstrated that when $\mathcal{X} = L^\infty$, any cash-subadditive, quasi-star-shaped risk measure is the pointwise minimum of a family of cash-subadditive, quasi-convex risk measures. The last thesis of Proposition 12 establishes that this assertion remains valid when we replace ‘quasi-star-shaped’ with ‘star-shaped’ and ‘quasi-convex’ with ‘convex’.

4 Representations à la Kusuoka

In this section, we first present quantile-based representations for star-shaped functionals that satisfy SSD- or CSD-consistency. That is, we extend the representations established in [10, 25, 29] to our setting. We assume that \mathcal{X} possesses a rearrangement-invariant structure. This assumption is satisfied by commonly used spaces such as Orlicz spaces, Orlicz hearts and Marcinkiewicz spaces. As emphasized in Section 2, within this framework, the concepts of $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity and the Fatou property are equivalent for any law-invariant convex functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$. Additionally, when f is also monotonic, continuity from below is equivalent to the previous two properties (see Proposition 2.5 in [4] for further details). We are ready to state the main result of this section.

Proposition 13. *Let \mathcal{X} be a rearrangement-invariant space. A functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is CSD-consistent and star-shaped if and only if it can be represented as:*

$$f(X) = \min_{\gamma \in \Gamma} \tilde{f}_\gamma(X) = \min_{\gamma \in \Gamma} \sup_{Y \in L^\infty} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{f}_\gamma^*(Y) \right\}, \quad X \in \mathcal{X}, \quad (4.1)$$

where $(\tilde{f}_\gamma)_{\gamma \in \Gamma}$ is the family of functionals provided in Corollary 8, and \tilde{f}_γ^* is given by:

$$\tilde{f}_\gamma^*(Y) = \sup_{X \in \mathcal{X}} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{f}_\gamma(X) \right\}, \quad Y \in \mathcal{X}^*, \gamma \in \Gamma.$$

In addition, if $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is an SSD-consistent and star-shaped risk measure, then the supremum in Equation (4.1) can be taken over L_+^∞ , obtaining:

$$\rho(X) = \min_{\gamma \in \Gamma} \tilde{\rho}_\gamma(X) = \min_{\gamma \in \Gamma} \sup_{Y \in L_+^\infty} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{\rho}_\gamma^*(Y) \right\}, \quad X \in \mathcal{X}.$$

Here, $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ is the family of risk measures provided in Theorem 7, and

$$\tilde{\rho}_\gamma^*(Y) = \sup_{X \in \mathcal{X}} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{\rho}_\gamma(X) \right\}, \quad Y \in \mathcal{X}^*, \gamma \in \Gamma.$$

Finally, $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a CSD-consistent and positively homogeneous functional if and only if the following representation holds:

$$f(X) = \min_{\gamma \in \Gamma} \tilde{f}_\gamma(X) = \min_{\gamma \in \Gamma} \sup_{Y \in \mathcal{M}^\gamma} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta \right\}, \quad X \in \mathcal{X},$$

where Γ is a set of indices and $\mathcal{M}^\gamma \subseteq L^\infty$ for each $\gamma \in \Gamma$. If f is also monotone (thus, it is a risk measure), then the family of sets $(\mathcal{M}^\gamma)_{\gamma \in \Gamma}$ can be chosen such that $\mathcal{M}^\gamma \subseteq L_+^\infty$.

Proof. The ‘if’ part is straightforward. Let us prove the ‘only if’ implication.

The CSD-consistent and star-shaped case

According to Proposition 5.1 in [4], if \tilde{f}_γ is law-invariant, convex and $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuous, it admits the representation:

$$\tilde{f}_\gamma(X) = \sup_{Y \in L^\infty} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{f}_\gamma^*(Y) \right\}, \quad X \in \mathcal{X},$$

valid for any $\gamma \in \Gamma$. Here, $\tilde{f}_\gamma^*(Y)$ is given by:

$$\tilde{f}_\gamma^*(Y) = \sup_{X \in \mathcal{X}} \left\{ \int_0^1 \text{VaR}_\beta(X) \text{VaR}_\beta(Y) d\beta - \tilde{f}_\gamma(X) \right\}, \quad Y \in \mathcal{X}^*.$$

Thus, it suffices to establish that \tilde{f}_γ is $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuous for any $\gamma \in \Gamma$, given that CSD-consistency and convexity of \tilde{f}_γ have already been shown in Corollary 8, recalling that CSD-consistency implies law-invariance (see Remark 5).

Equivalent expression for \tilde{f}_Z : Referring back to Corollary 8 and considering the explicit form of the law-invariant convex functional \tilde{f}_γ , where $\gamma := Z \in \text{Dom}(f)$, involved in the representation of f (assuming for brevity that $f(0) = 0$), we can express \tilde{f}_Z as follows:

$$\tilde{f}_Z(X) = \begin{cases} \inf_{\alpha \in A_X} \{\alpha f(Z)\} & \text{if } A_X := \{\alpha \in [0, 1] : \exists Y \succeq_c X, Y \sim \alpha Z\} \neq \emptyset, \\ +\infty & \text{otherwise.} \end{cases}$$

We also recall the definition of $(f_Z)_{Z \in \Gamma}$:

$$f_Z(X) = \begin{cases} \alpha f(Z) & \text{if } \exists \alpha \in (0, 1] \text{ s.t. } X \sim \alpha Z, X \not\sim 0 \\ 0 & \text{if } X \sim 0 \\ +\infty & \text{otherwise.} \end{cases}$$

Note that for any fixed $X \in \mathcal{X}$ the infimum in the definition of $\tilde{f}_Z(X)$ is always attained as soon as $A_X \neq \emptyset$. Indeed, by definition of the infimum, there exist a sequence of real numbers $(\alpha_j)_{j \in \mathbb{N}} \subseteq [0, 1]$ and a sequence of random variables $(Y_j)_{j \in \mathbb{N}}$ verifying $Y_j \sim \alpha_j Z$ and $Y_j \succeq_c X$ for all $j \in \mathbb{N}$ such that $\lim_{j \rightarrow \infty} \alpha_j f(Z) = \inf_{\alpha \in A_X} \{\alpha f(Z)\}$. We assume $f(Z) \neq 0$, otherwise the statement is trivial, once observed that any $\alpha \in A_X$ is a minimizer in this case. In particular, we have that

$$\lim_{j \rightarrow \infty} \alpha_j = \frac{\inf_{\alpha \in A_X} \{\alpha f(Z)\}}{f(Z)} \in [0, 1].$$

We want to prove that $\bar{\alpha} := \lim_{j \rightarrow \infty} \alpha_j$ is actually a minimizer. Let us consider the random variable $\bar{Y} := \bar{\alpha}Z$. We need to check that $\bar{Y} \succeq_c X$ and $\bar{Y} \sim \bar{\alpha}Z$. The second condition is true by construction while the first condition follows from positive homogeneity of Expected Shortfall: $ES_\beta(Y_j) = \alpha_j ES_\beta(Z) \geq ES_\beta(X)$ for all $j \in \mathbb{N}$, and letting $j \rightarrow \infty$ we obtain $ES_\beta(\bar{Y}) = ES_\beta(\bar{\alpha}Z) = \lim_{j \rightarrow \infty} \alpha_j ES_\beta(Z) \geq ES_\beta(X)$ for all $\beta \in [0, 1]$. In addition, it results that

$$\lim_{j \rightarrow \infty} \mathbb{E}(Y_j) = \lim_{j \rightarrow \infty} \alpha_j \mathbb{E}(Z) = \mathbb{E}(\bar{\alpha}Z) = \mathbb{E}(Y) = \mathbb{E}(X),$$

thus the thesis follows.

Lebesgue property of ES_β : We claim that ES_β possesses the Lebesgue property for any $\beta \in [0, 1]$. Based on Proposition 2.35 in [38], for any $X, Y \in L^1$ it follows that:

$$|ES_\beta(X) - ES_\beta(Y)| \leq \frac{1}{1 - \beta} \|X - Y\|_1.$$

Given the hypotheses, $\sup_{n \in \mathbb{N}} |X_n| \in \mathcal{X} \subseteq L^1$, $X \in \mathcal{X} \subseteq L^1$, and $X_n \rightarrow X$ a.s., the dominated convergence theorem implies $X_n \rightarrow X$ in L^1 . Consequently,

$$|ES_\beta(X_n) - ES_\beta(X)| \leq \frac{1}{1 - \beta} \|X_n - X\|_1 \rightarrow 0,$$

yielding $ES_\beta(X_n) \rightarrow ES_\beta(X)$ for any $\beta \in [0, 1]$.

$\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity of \tilde{f}_Z : As \mathcal{X} is a rearrangement-invariant space, demonstrating the Fatou property of \tilde{f}_γ for any $\gamma \in \Gamma$ is sufficient. This property requires showing that for a sequence $(X_n)_{n \in \mathbb{N}} \subseteq \mathcal{X}$ and $X \in \mathcal{X}$ with $X_n \rightarrow X$ a.s. and $\sup_{n \in \mathbb{N}} |X_n| \in \mathcal{X}$, it holds that $\liminf_{n \in \mathbb{N}} \tilde{f}_\gamma(X_n) \geq \tilde{f}_\gamma(X)$. Based on the preceding argument and assuming that $\tilde{f}_Z(X) \neq +\infty$, it is evident that $\liminf_{n \in \mathbb{N}} \tilde{f}(X_n) \neq +\infty$ only if there are infinitely many $n \in \mathbb{N}$ for which there exists $Y_n \succeq_c X_n$ with $Y_n \sim \alpha_n Z$, $\alpha_n \in [0, 1]$ and $\tilde{f}(X_n) = f_Z(Y_n) = \alpha_n f(Z)$. This allows focusing on such sequences. We need to verify that $\liminf_{n \rightarrow \infty} \tilde{f}_Z(X_n) \geq \tilde{f}_Z(X)$. It holds that:

$$\liminf_{n \rightarrow \infty} \tilde{f}_Z(X_n) = \liminf_{n \rightarrow \infty} f_Z(Y_n) = \liminf_{n \rightarrow \infty} \alpha_n f(Z) = \tilde{\alpha} f(Z) = f_Z(\tilde{Y}),$$

where $\tilde{Y} := \tilde{\alpha} Z$ with $\tilde{\alpha} := \liminf_{n \rightarrow \infty} \alpha_n$. The last equality results from observing that $\tilde{\alpha} = \liminf_{n \rightarrow \infty} \alpha_n \in [0, 1]$ due to the fact that $\alpha_n \in [0, 1]$ for all $n \in \mathbb{N}$, and in accordance with the definition of f_Z . If we show that $\tilde{Y} \succeq_c X$ (thus, $\tilde{\alpha} \in A_X$), then we can conclude $f_Z(\tilde{Y}) \geq \inf_{\alpha \in A_X} \alpha f(Z) = \tilde{f}_Z(X)$, leading to $\liminf_{n \rightarrow \infty} \tilde{f}_Z(X_n) \geq \tilde{f}_Z(X)$. To establish this, we need to verify $ES_\beta(\tilde{Y}) \geq ES_\beta(X)$ and $\mathbb{E}(\tilde{Y}) = \mathbb{E}(X)$. We have:

$$\begin{aligned} ES_\beta(\tilde{Y}) &= \tilde{\alpha} ES_\beta(Z) = \liminf_{n \rightarrow \infty} \alpha_n ES_\beta(Z) \\ &= \liminf_{n \rightarrow \infty} ES_\beta(Y_n) \geq \liminf_{n \rightarrow \infty} ES_\beta(X_n) = ES_\beta(X), \end{aligned}$$

where the inequality is due to $Y_n \succeq_c X_n$ and the last equality holds by the Lebesgue property of ES_β . Thus, $ES_\beta(\tilde{Y}) \geq ES_\beta(X)$ for any $\beta \in [0, 1]$. Analogously, by the dominated convergence theorem, we have

$$\mathbb{E}(\tilde{Y}) = \liminf_{n \rightarrow \infty} \mathbb{E}(Y_n) = \liminf_{n \rightarrow \infty} \mathbb{E}(X_n) = \mathbb{E}(X).$$

By a similar argument involving contradiction, it can be established that when $\tilde{f}_\gamma(X) = +\infty$, $\liminf_{n \rightarrow \infty} \tilde{f}(X_n) = +\infty$, hence \tilde{f}_Z verifies the Fatou property, yielding the $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity.

The SSD-consistent and star-shaped case

Let us consider an SSD-consistent and star-shaped risk measure $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$. For brevity we suppose $\rho(0) = 0$. Our goal is to establish the $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity of $\tilde{\rho}_\gamma$ for any $\gamma \in \Gamma$. Importantly, the SSD-consistency of ρ inherently includes both monotonicity and law-invariance. Consequently, we can apply Proposition 5.1 as presented in [4] once more to derive the desired dual representation for $\tilde{\rho}_\gamma$.

Equivalent expression for $\tilde{\rho}_Z$: By Theorem 7, we know that $\tilde{\rho}_\gamma$ takes the following form, with $\gamma := Z \in \text{Dom}(f)$:

$$\tilde{\rho}_Z(X) := \inf\{\rho_Z(Y) : Y \succeq_2 X\},$$

which can be written as:

$$\tilde{\rho}_Z(X) = \begin{cases} \inf_{\alpha \in A_X} \{\alpha \rho(Z)\} & \text{if } A_X := \{\alpha \in [0, 1] : \exists Y \succeq_2 X, Y \sim \alpha Z\} \neq \emptyset, \\ +\infty & \text{otherwise,} \end{cases} \quad (4.2)$$

where

$$\rho_Z(X) = \begin{cases} \alpha \rho(Z) & \text{if } \exists \alpha \in (0, 1] \text{ s.t. } X \sim \alpha Z, X \not\sim 0 \\ 0 & \text{if } X \sim 0 \\ +\infty & \text{otherwise.} \end{cases}$$

Note that for any fixed $X \in \mathcal{X}$ the infimum in the definition of $\tilde{\rho}_Z(X)$ is always attained as soon as $A_X \neq \emptyset$. The proof of this statement is analogous to the case of CSD-consistency.

$\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity of $\tilde{\rho}_Z$: Under the monotonicity assumption, $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity is equivalent to continuity from below. Let $(X_n)_{n \in \mathbb{N}} \subseteq \mathcal{X}$ be a sequence such that $X_n \uparrow X \in \mathcal{X}$ a.s.; then we want to show that $\tilde{\rho}_\gamma(X_n) \rightarrow \tilde{\rho}_\gamma(X)$ for any $\gamma \in \Gamma$. By monotonicity, it is enough to prove that $\liminf_{n \rightarrow \infty} \tilde{\rho}_\gamma(X_n) \geq \tilde{\rho}_\gamma(X)$ for any $\gamma \in \Gamma$. By hypotheses we have $X_n \uparrow X$, hence the continuity from below of Expected Shortfall ensures that $ES_\beta(X) = \lim_{n \rightarrow \infty} ES_\beta(X_n)$. We start assuming $\tilde{\rho}_Z(X) < +\infty$. Once again, we can consider $(X_n)_{n \in \mathbb{N}}$ such that for each $n \in \mathbb{N}$ there exists $Y_n \succeq_2 X_n$ verifying $Y_n \sim \alpha_n Z$ with $\alpha_n \in [0, 1]$ attaining the minimum of the set A_{X_n} . Thus,

$$ES_\beta(X) = \lim_{n \rightarrow \infty} ES_\beta(X_n) \leq \liminf_{n \rightarrow \infty} ES_\beta(Y_n) = \tilde{\alpha} ES_\beta(Z),$$

for any $\beta \in [0, 1)$, with $\tilde{\alpha} := \liminf_{n \rightarrow \infty} \alpha_n$, where the inequality follows from the SSD-consistency of ES_β and the relation $Y_n \succeq_2 X_n$ for all $n \in \mathbb{N}$. The above chain of inequalities, the definition of $\tilde{\rho}_Z$, the SSD-consistency of $\tilde{\rho}_Z$ and the relation $\tilde{\rho}_Z \leq \rho_Z$ yield:

$$\tilde{\rho}_Z(X) \leq \liminf_{n \rightarrow \infty} \tilde{\rho}_Z(Y_n) \leq \liminf_{n \rightarrow \infty} \rho_Z(\alpha_n Z) = \tilde{\alpha} \rho(Z) = \liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n),$$

hence $\liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) \geq \rho(X)$. Similarly, if we assume $\rho_Z(X) = +\infty$ it can be proved that $\liminf_{n \rightarrow \infty} \rho_Z(X_n) = +\infty$, thus the thesis follows.

The positively homogeneous case: In the case of positive homogeneity, we can similarly deduce the $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity of $\tilde{\rho}_Z$ and \tilde{f}_Z for SSD- and CSD-consistent functionals, respectively. The key difference is that in this setting, the parameter α falls within the interval $\alpha \in [0, +\infty)$ rather than the previously considered $\alpha \in [0, 1]$. \square

The following corollary shows that we are also able to characterize cash-additive and SSD-consistent star-shaped risk measures through a representation à la Kusuoka, as robustification of Expected Shortfall. These findings extend Equation (3.9) of [35] to star-shapedness and to a broader space than L^∞ .

Corollary 14. *Let \mathcal{X} be a rearrangement-invariant space. A risk measure $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is SSD-consistent, star-shaped and cash-additive if and only if ρ admits the representation:*

$$\rho(X) = \min_{\gamma \in \Gamma} \tilde{\rho}_\gamma(X) = \min_{\gamma \in \Gamma} \sup_{\mu \in \mathcal{P}([0,1])} \left\{ \int_0^1 ES_s(X) d\mu(s) - \alpha_\gamma(\mu) \right\},$$

where $\mathcal{P}([0,1])$ is the set of probability measures on $([0,1], \mathcal{B}([0,1]))$, Γ is a set of indices, $(\tilde{\rho}_\gamma)_{\gamma \in \Gamma}$ is the family of risk measures provided in Proposition 9 and α_γ is a penalty function defined for any $\mu \in \mathcal{P}([0,1])$ as:

$$\alpha_\gamma(\mu) := \sup \left\{ \int_0^1 ES_s(X) d\mu(s); X \in L^\infty, \rho_\gamma(X) \geq 0 \right\}.$$

Proof. The ‘if’ part is trivial. Let us prove only the converse implication. The proof is similar to the proof of Proposition 13. For brevity we suppose $\rho(0) = 0$ and we use the same notation as in Proposition 13. We recall that for any $\gamma := Z \in \text{Dom}(\rho)$ the functional $\tilde{\rho}_Z$ is given by

$$\tilde{\rho}_Z(X) := \inf \{ \rho_Z(Y) : Y \succeq_2 X \}, X \in \mathcal{X}.$$

We know that $\tilde{\rho}_Z$ is SSD-consistent, convex, monotone and cash-additive. If we can prove that $\tilde{\rho}_Z$ is also continuous from below, then the thesis follows from Proposition 5.12 in [4], as law-invariance is implied by SSD-consistency and $\sigma(\mathcal{X}, \mathcal{X}^*)$ -lower semicontinuity is equivalent to continuity from below for a monotone functional defined on a rearrangement invariant space.

Equivalent definition of $\tilde{\rho}_Z$: Once again we can write an equivalent expression for $\tilde{\rho}_Z$. Fixing $Z \in \text{Dom}(\rho)$, for each $X \in \mathcal{X}$ we define the set $B_X := \{(\alpha, c) \in [0, 1] \times \mathbb{R} : \exists Y \in \mathcal{X} \text{ s.t. } Y \succeq_2 X \text{ and } Y \sim \alpha Z + c\}$. Then ρ_Z takes the following form:

$$\tilde{\rho}_Z(X) = \begin{cases} \inf_{(\alpha, c) \in B_X} \{\alpha \rho(Z) + c\} & \text{if } B_X \neq \emptyset, \\ +\infty & \text{otherwise.} \end{cases}$$

We observe that when B_X is non-empty the infimum in the previous equation must be finite given that $\tilde{\rho}_Z(X) \geq \rho(X) > -\infty$. In addition, when B_X is non-empty the infimum is in fact a minimum. Indeed, arguing as in the proof of Proposition 13, there exist a sequence of real numbers $(\alpha_j, c_j)_{j \in \mathbb{N}} \subseteq [0, 1] \times \mathbb{R}$ and a sequence of random variables Y_j such that $\alpha_j \rho(Z) + c_j \rightarrow \inf_{(\alpha, c) \in B_X} \{\alpha \rho(Z) + c\}$ for $j \rightarrow \infty$ and $Y_j \succeq X, Y_j \sim \alpha_j Z + c_j$ for any $j \in \mathbb{N}$. By compactness of $[0, 1]$ there exists a sub-sequence $(\alpha_{j_i})_{i \in \mathbb{N}}$ converging to some $\bar{\alpha} \in [0, 1]$. Moreover, $\lim_{i \rightarrow \infty} \{\alpha_{j_i} \rho(Z) + c_{j_i}\} = \inf_{(\alpha, c) \in B_X} \{\alpha \rho(Z) + c\}$, hence $\lim_{i \rightarrow \infty} c_{j_i}$ exists and we define $\bar{c} := \lim_{i \rightarrow \infty} c_{j_i}$. Let us consider $\bar{Y} := \bar{\alpha} Z + \bar{c}$. By construction we have $\bar{Y} \sim \bar{\alpha} Z + \bar{c}$. Moreover, by cash-additivity and positive homogeneity of Expected Shortfall we have $\alpha_{j_i} ES_\beta(Z) + c_{j_i} \geq X$ for any $\beta \in [0, 1)$ and $i \in \mathbb{N}$. Letting $i \rightarrow \infty$ we get $ES_\beta(\bar{Y}) \geq X$ for any $\beta \in [0, 1)$, hence the infimum is attained at $(\bar{\alpha}, \bar{c}) \in [0, 1] \times \mathbb{R}$.

Continuity from below of $\tilde{\rho}_Z$: Given $X_n \uparrow X$ we want to prove that $\lim_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) = \tilde{\rho}_Z(X)$. Due to monotonicity, it suffices to demonstrate $\liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) \geq \tilde{\rho}_Z(X)$. We assume $\tilde{\rho}_Z(X) < +\infty$, which, by monotonicity, implies $\liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) < +\infty$. Thus, we can consider a sequence $(X_n)_{n \in \mathbb{N}}$ such that $\tilde{\rho}_Z(X_n) < +\infty \quad \forall n \in \mathbb{N}$, verifying $\tilde{\rho}_Z(X_n) = \rho_Z(Y_n) = \alpha_n \rho(Z) + c_n$. Here, for each $n \in \mathbb{N}$, $(\alpha_n, c_n) \in [0, 1] \times \mathbb{R}$ attains the infimum of B_{X_n} , with $Y_n \succeq_2 X_n$ and $Y_n \sim \alpha_n Z + c_n$. Because $[0, 1]$ is a compact set, we can extract a sub-sequence $(\alpha_{n_k})_{k \in \mathbb{N}}$ such that $\alpha_{n_k} \rightarrow \tilde{\alpha} \in [0, 1]$. By monotonicity of $\tilde{\rho}_Z$, we infer that $\tilde{\rho}_Z(X_n)$ must converge to some $l \in \mathbb{R}$ (otherwise, we would have $\liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) = +\infty$, which contradicts the hypothesis). Thus, also $\tilde{\rho}_Z(X_{n_k}) \rightarrow l$ and then $c_{n_k} = \tilde{\rho}_Z(X_{n_k}) - \alpha_{n_k} \rho(Z) \rightarrow l - \tilde{\alpha} \rho(Z) := \tilde{c} \in \mathbb{R}$.

Since $X_n \uparrow X$, continuity from below of Expected Shortfall ensures that $ES_\beta(X) = \lim_{n \rightarrow \infty} ES_\beta(X_n)$. Thus,

$$ES_\beta(X) = \lim_{k \rightarrow \infty} ES_\beta(X_{n_k}) \leq \liminf_{k \rightarrow \infty} ES_\beta(Y_{n_k}) = \tilde{\alpha} ES_\beta(Z) + \tilde{c},$$

for any $\beta \in [0, 1)$, where the inequality follows from the SSD-consistency of ES_β and the relation $Y_{n_k} \succeq_2 X_{n_k}$ for all $k \in \mathbb{N}$. The above chain of inequalities, the definition of $\tilde{\rho}_Z$ and the SSD-consistency of $\tilde{\rho}_Z$ yield:

$$\begin{aligned} \tilde{\rho}_Z(X) &\stackrel{\text{SSD}}{\leq} \lim_{k \rightarrow \infty} \tilde{\rho}_Z(Y_{n_k}) \stackrel{\tilde{\rho}_Z \leq \rho_Z}{\leq} \lim_{k \rightarrow \infty} \rho_Z(Y_{n_k}) \\ &\stackrel{\text{def. } \rho_Z}{=} \tilde{\alpha} \rho(Z) + \tilde{c} = \lim_{k \rightarrow \infty} \{\alpha_{n_k} \rho(Z) + c_{n_k}\} \\ &= \lim_{k \rightarrow \infty} \tilde{\rho}_Z(X_{n_k}) = \liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n). \end{aligned}$$

If $\tilde{\rho}_Z(X) = +\infty$ we can proceed by contradiction, proving that also $\liminf_{n \rightarrow \infty} \tilde{\rho}_Z(X_n) = +\infty$. The thesis follows. \square

5 Law-Invariant Star-Shaped Functionals as Robustification of Value-at-Risk

While we have proved that every SSD-consistent (resp. CSD-consistent) star-shaped functional arises as the minimum of SSD-consistent (resp. CSD-consistent) convex functionals, we

aim to characterize the bigger family of law-invariant and star-shaped functionals in terms of Value-at-Risk. Indeed, as explained in the previous sections, not every law-invariant star-shaped functional can be described as the minimum of a family of law-invariant convex functionals, having the Value-at-Risk as a classical counter-example to this statement. See also Section 7 in [9] and the references therein for a thorough discussion on this topic. More specifically, we are seeking to establish representations in a similar spirit to those provided in Theorem 5 of [9] and Proposition A.4 of [28]. In [9], law-invariant star-shaped monetary risk measures appear as a robustification of Value-at-Risk, where the generalized scenarios are represented by a penalty function dependent on the β -level of VaR_β . A similar representation is found in [28] for cash-subadditive risk measures, even when star-shapedness is not a requirement in this case. In the following proposition we show that this representation can be extended to the more general setting in which cash-additivity (or cash-subadditivity) and monotonicity are discarded, and \mathcal{X} is a (much) more general space than L^∞ . The following results are presented under the assumption of normalization. It is important to note that the proof strategy remains applicable even for a *non-normalized* functional.

Proposition 15. *Let $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a functional such that $f(0) = 0$. Then, f is law-invariant and star-shaped if and only if it admits the following representation:*

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}, \quad X \in \mathcal{X}. \quad (5.1)$$

Here, for each $X \in \mathcal{X}$, Γ_X is a set of random variables such that $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0,1]$. In addition, for any $X, X' \in \mathcal{X}$ such that $X \sim X'$, it holds that $\Gamma_X = \Gamma_{X'}$. For each $X \in \mathcal{X}$ and $Z \in \Gamma_X$, $\gamma_Z^X : (0,1) \rightarrow \mathbb{R} \cup \{-\infty\}$ is an increasing function such that $\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0$ and for any $\lambda \in (0,1], Z \in \Gamma_X$ it holds that $\gamma_{\lambda Z}^{\lambda X} \geq \lambda \gamma_Z^X$.

Furthermore, for each $X \in \mathcal{X}$, the set Γ_X takes the following form:

$$\Gamma_X := \{Z \in \text{Dom}(f) : X \sim \alpha Z, \alpha \in [0,1]\},$$

and the function $\gamma_Z^X : (0,1) \rightarrow \mathbb{R} \cup \{-\infty\}$ with $Z \in \Gamma_X$ is defined by the formula:

$$\gamma_Z^X(\beta) = \alpha(VaR_\beta(Z) - f(Z)),$$

with the convention that if $\Gamma_X = \emptyset$ then $\gamma_Z^X = -\infty$.

With the same notation as above, $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is law-invariant and positively homogeneous if and only if f can be represented as in Equation (5.1), with $\lambda \in [0,+\infty)$, $\lambda\Gamma_X = \Gamma_{\lambda X}$ and $\gamma_{\lambda Z}^{\lambda X} = \lambda \gamma_Z^X$. In addition, the set Γ_X can be defined for all $\alpha \in [0,+\infty)$.

Proof. We prove the statement in the star-shapedness case, the positively homogeneous case follows similarly. We assume $X \in \text{Dom}(f)$, otherwise the statements are trivial.⁸

We start by proving the ‘only if’ part. It is worth noting that Γ_X is well defined, given that if $X \sim \alpha Z$, with $\alpha \in [0,1]$ or $\alpha \in [0,+\infty)$, then α is unique, as we have shown in the first part of the proof of Theorem 7.

Min-sup representation: Let us consider the functional

$$X \mapsto \tilde{\gamma}_{Z,\beta}(X) := VaR_\beta(X) - \gamma_Z^X(\beta),$$

with $\beta \in (0,1)$, and $Z \in \Gamma_X$. It holds that:

$$\begin{aligned} \tilde{\gamma}_{Z,\beta}(X) &= VaR_\beta(X) - \alpha(VaR_\beta(Z) - f(Z)) \\ &= VaR_\beta(X) - VaR_\beta(\alpha Z) + \alpha f(Z) \\ &= \alpha f(Z) \geq f(\alpha Z) = f(X), \end{aligned}$$

⁸It is worth noting that $\Gamma_X \neq \emptyset \iff X \in \text{Dom}(f)$. Indeed, if $X \in \text{Dom}(f)$, then we can choose $Z = X$, which clearly lies in Γ_X with $\alpha = 1$. Conversely, if $X \notin \text{Dom}(f)$, then for any $Z \in \mathcal{X}$ such that $X \sim \alpha Z$, we have that $\alpha f(Z) \geq f(\alpha Z) = f(X) = +\infty$. Thus, $f(Z) = +\infty$, and hence $Z \notin \Gamma_X$ since $Z \notin \text{Dom}(f)$. This observation applies, with minor modifications, to all the results presented in Section 5.

where law-invariance of VaR implies $VaR_\beta(X) - VaR_\beta(\alpha Z) = 0$, while the inequality is due to star-shapedness of f and the last equality follows from law-invariance of f . Thus, $\inf_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \tilde{\gamma}_{Z,\beta}(X) \geq f(X)$. Observing that $X \in \Gamma_X$ and $\tilde{\gamma}_{X,\beta}(X) = f(X)$ for any $\beta \in (0,1)$ we get:

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}.$$

Properties of Γ_X : It is a routine verification to prove that $\Gamma_{\lambda X} \subseteq \lambda \Gamma_X$ for any $\lambda \in (0,1]$, once observed that $Z \in \text{Dom}(f) \implies \lambda Z \in \text{Dom}(f)$. Indeed, if $Z \in \text{Dom}(f)$ then $f(Z) < +\infty$ and using star-shapedness of f it follows that $f(\lambda Z) \leq \lambda f(Z) < +\infty$, thus $\lambda Z \in \text{Dom}(f)$. We only need to demonstrate that if $X \sim X'$, then $\Gamma_X = \Gamma_{X'}$. Indeed, if $X \sim X'$, it also follows that $X \sim X'$. Consequently, we have:

$$Z \in \Gamma_X \iff \exists \alpha \in [0,1] \mid X \sim \alpha Z \iff X' \sim \alpha Z \iff Z \in \Gamma_{X'}.$$

Properties of γ_Z^X : Let us fix $\lambda \in (0,1]$. Due to the star-shaped property of ρ and the set inclusion $\lambda \Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0,1]$, it follows that for any $Z \in \Gamma_X$ and $\beta \in (0,1)$:

$$\lambda \gamma_Z^X = \lambda \alpha (VaR_\beta(Z) - f(Z)) \leq \alpha (VaR_\beta(\lambda Z) - f(\lambda Z)) = \gamma_{\lambda Z}^{\lambda X}.$$

In addition, we have:

$$\rho(0) = \min_{Z \in \Gamma_0} \sup_{\beta \in (0,1)} \{-\gamma_Z^0(\beta)\} = \min_{Z \in \Gamma_0} \{-\gamma_Z^0(0^+)\} = 0,$$

where the second equality follows from increasing monotonicity of VaR_β w.r.t. β , while the last equality is due to normalization. Thus, $\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0$.

Now we are ready to prove the converse implication. We want to show that

$$\rho(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}, \quad X \in \mathcal{X},$$

is normalized, star-shaped and law-invariant.

Normalization: We have that:

$$\rho(0) = \min_{Z \in \Gamma_0} \sup_{\beta \in (0,1)} \{-\gamma_Z^0(\beta)\} = -\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0.$$

Star-shapedness: Let us consider $\lambda \in (0,1]$ and $X \in \mathcal{X}$. We have the following inequalities:

$$\begin{aligned} \rho(\lambda X) &= \min_{Z \in \Gamma_{\lambda X}} \sup_{\beta \in (0,1)} \{VaR_\beta(\lambda X) - \gamma_Z^{\lambda X}(\beta)\} \\ &= \lambda \min_{Z \in \Gamma_{\lambda X}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \frac{1}{\lambda} \gamma_Z^{\lambda X}(\beta)\} \\ &\leq \lambda \min_{Z \in \lambda \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \frac{1}{\lambda} \gamma_Z^{\lambda X}(\beta)\} \\ &\leq \lambda \min_{Z \in \lambda \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_{\frac{Z}{\lambda}}^X(\beta)\} \\ &= \lambda \min_{Z' \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_{Z'}^X(\beta)\} = \lambda \rho(X), \end{aligned}$$

where the inequalities follow from $\gamma_{\frac{Z}{\lambda}}^X \geq \lambda \gamma_Z^{\lambda X}$ and $\lambda \Gamma_X \subseteq \Gamma_{\lambda X}$.

Law-invariance: This assertion follows from the law-invariance of VaR and the relation $\Gamma_X = \Gamma_{X'}$ as long as $X \sim X'$. Thus, we can express it as follows:

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\} = \min_{Z \in \Gamma_{X'}} \sup_{\beta \in (0,1)} \{VaR_\beta(X') - \gamma_Z^X(\beta)\} = f(X').$$

□

Corollary 16. *A functional $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is normalized, law-invariant, star-shaped and cash-additive if and only if it admits the representation:*

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\},$$

where for each $X \in \mathcal{X}$, Γ_X is a set of random variables such that $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0, 1]$, $\Gamma_X = \Gamma_{X'}$ whenever $X \sim X'$ and $\Gamma_{X+m} = \Gamma_X$ for all $m \in \mathbb{R}$. In addition, $\gamma_Z^X : (0, 1) \rightarrow \mathbb{R} \cup \{-\infty\}$ is an increasing function such that $\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0$. Moreover, for any $Z \in \Gamma_X$, $\lambda \in (0, 1]$ it holds that $\lambda\gamma_Z^X = \gamma_{\lambda Z}^{\lambda X}$ and for any $m \in \mathbb{R}$ it follows that $\gamma_Z^{X+m} = \gamma_Z^X$.

Furthermore, for each $X \in \mathcal{X}$, the set Γ_X takes the following form:

$$\Gamma_X := \{Z \in \text{Dom}(f) : X \sim \alpha Z + c, (\alpha, c) \in [0, 1] \times \mathbb{R}, f(Z) \leq 0\},$$

with $(\alpha, c) = (0, X)$ if X is constant. The function $\gamma_Z^X : (0, 1) \rightarrow \mathbb{R} \cup \{-\infty\}$ with $Z \in \Gamma_X$ is defined by the formula:

$$\gamma_Z^X(\beta) = \alpha VaR_\beta(Z),$$

with that convention that if $\Gamma_X = \emptyset$ then $\gamma_Z^X = -\infty$.

With the same notation as above, $f : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is law-invariant, positively homogeneous and cash-additive if and only if f can be represented as in Equation (5.1), with $\lambda \in [0, +\infty)$ and $\lambda\Gamma_X = \Gamma_{\lambda X}$. In addition, the set Γ_X can be defined for any $\alpha \in [0, +\infty)$.

Proof. We prove the statement in the star-shapedness case, the positively homogeneous case follows similarly. We commence by establishing the ‘only if’ part. It is important to note that for any $Z \in \Gamma_X$, the function γ_Z^X is well-defined. Indeed, if X is non-constant and $X \sim \alpha Z + c$ with $(\alpha, c) \in [0, 1] \times \mathbb{R}$, then, as demonstrated in the first part of the proof of Proposition 9, (α, c) is a unique pair of values. Additionally, if X is constant, we have $\gamma_Z^X \equiv 0$, for any $Z \in \Gamma_X = \{Z \in \text{Dom}(f) : f(Z) \leq 0\}$. Henceforth, we assume $X \in \text{Dom}(f)$ with X non-constant, otherwise the statements are trivial.

Min-sup representation: Let us fix $X \in \text{Dom}(f)$, $\beta \in (0, 1)$, and $Z \in \Gamma_X$. It holds that:

$$\begin{aligned} \tilde{\gamma}_{Z,\beta} &:= VaR_\beta(X) - \alpha(VaR_\beta(Z) - f(Z)) \\ &= VaR_\beta(X) - VaR_\beta(\alpha Z + c) + \alpha f(Z) + c \\ &= \alpha f(Z) + c \geq f(\alpha Z + c) = f(X), \end{aligned}$$

where law-invariance and cash-additivity of VaR implies $VaR_\beta(X) - VaR_\beta(\alpha Z + c) = 0$, while the inequality is due to star-shapedness and cash-additivity of f and the last equality follows from law-invariance of f . Thus,

$$\inf_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha(VaR_\beta(Z) - f(Z))\} \geq f(X).$$

Since cash-additivity implies $f(X - f(X)) = 0$ we have $X - f(X) \in \Gamma_X$, with $(\alpha, c) = (1, f(X))$ and $\tilde{\gamma}_{X,\beta}(X) = f(X)$. Thus,

$$\begin{aligned} f(X) &= \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha(VaR_\beta(Z) - f(Z))\} \\ &= \min_{\substack{Z \in \Gamma_X \\ f(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha VaR_\beta(Z)\} \\ &= \min_{\substack{Z \in \Gamma_X \\ f(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}. \end{aligned}$$

Furthermore, given that $Z \in \Gamma_X \implies f(Z) \leq 0$ it follows that:

$$\begin{aligned} f(X) &= \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha(VaR_\beta(Z) - f(Z))\} \\ &\leq \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha VaR_\beta(Z)\} \\ &\leq \min_{\substack{Z \in \Gamma_X \\ f(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \alpha VaR_\beta(Z)\} = f(X), \end{aligned}$$

hence

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}.$$

Properties of Γ_X : Law-invariance of Γ_X can be proved similarly as in Proposition 15. We now prove the inclusion $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0, 1]$. Let $Z \in \lambda\Gamma_X$. Then there exists $Z' \in \Gamma_X$ such that $Z = \lambda Z'$ and $X = \alpha Z' + c$ with $f(Z') \leq 0$ and $(\alpha, c) \in [0, 1] \times \mathbb{R}$. It holds that:

$$\lambda X \sim \lambda(\alpha Z' + c) = \alpha Z + c',$$

where $c' := \lambda c \in \mathbb{R}$. In addition, we have $f(Z) = f(\lambda Z') \leq \lambda f(Z') \leq 0$, thus $Z \in \Gamma_{\lambda X}$. It is worth noting that the converse inclusion $\Gamma_{\lambda X} \subseteq \lambda\Gamma_X$ does not hold in general. In addition, for $m \in \mathbb{R}$ and $X \in \mathcal{X}$ we have that $Z \in \Gamma_X \iff Z \in \Gamma_{X+m}$. Indeed, if $Z \in \Gamma_X$ then there exist $(\alpha, c) \in [0, 1] \times \mathbb{R}$ such that $X \sim \alpha Z + c$. Considering $c' := c + m$ we have $X + m \sim \alpha Z + c'$, thus $Z \in \Gamma_{X+m}$. So we showed that $\Gamma_X \subseteq \Gamma_{X+m}$; the converse inclusion can be proved analogously.

Properties of γ_Z^X : Let us fix $\lambda \in (0, 1]$. By the inclusion $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0, 1]$, it follows for any $Z \in \Gamma_X$ and $\beta \in (0, 1)$ that:

$$\lambda \gamma_Z^X = \lambda \alpha VaR_\beta(Z) = \alpha VaR_\beta(\lambda Z) = \gamma_{\lambda Z}^{\lambda X},$$

where last equality follows from $\lambda Z \in \lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ with $\lambda X = \alpha \lambda Z + \lambda c$. The equality $\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0$ follows as in Proposition 15. Finally, consider $m \in \mathbb{R}$. We have:

$$\gamma_Z^{X+m} = \alpha VaR(Z) = \gamma_Z^X,$$

where $\alpha \in [0, 1]$ is the same in the representation $X \sim \alpha Z + c$, whether Z is regarded as an element of Γ_X or as an element of Γ_{X+m} , as demonstrated earlier in the proof.

Now we are ready to prove the converse implication. We want to show that

$$f(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}, \quad X \in \mathcal{X},$$

is normalized, star-shaped, law-invariant and cash-additive. Normalization and law-invariance follows as in the proof of Proposition 15.

Star-shapedness: Let us consider $\lambda \in (0, 1]$. We have the following inequalities:

$$\begin{aligned} f(\lambda X) &= \min_{Z \in \Gamma_{\lambda X}} \sup_{\beta \in (0,1)} \{VaR_\beta(\lambda X) - \gamma_Z^{\lambda X}(\beta)\} = \\ &= \lambda \min_{Z \in \Gamma_{\lambda X}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \frac{1}{\lambda} \gamma_Z^{\lambda X}(\beta)\} \\ &= \lambda \min_{\lambda Z' \in \Gamma_{\lambda X}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \frac{1}{\lambda} \gamma_{\lambda Z'}^{\lambda X}(\beta)\} \\ &\leq \lambda \min_{\lambda Z' \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \frac{1}{\lambda} \gamma_{\lambda Z'}^{\lambda X}(\beta)\} \\ &= \lambda \min_{Z' \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_{Z'}^X(\beta)\} = \lambda f(X), \end{aligned}$$

where the inequality follows from $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$, while $\gamma_{\lambda Z}^{\lambda X} = \lambda\gamma_Z^X$ for any $Z \in \Gamma_X$ leads to the last equality.

Cash-additivity: Let $m \in \mathbb{R}$. We have:

$$\begin{aligned} f(X+m) &= \min_{Z \in \Gamma_{X+m}} \sup_{\beta \in (0,1)} \{VaR_\beta(X+m) - \gamma_Z^{X+m}(\beta)\} \\ &= \min_{Z \in \Gamma_{X+m}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^{X+m}(\beta)\} + m \\ &= \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\} + m = f(X) + m. \end{aligned}$$

The last equality follows from $\Gamma_{X+m} = \Gamma_X$ and $\gamma_Z^{X+m} = \gamma_Z^X$. \square

In the subsequent lemma, we introduce a method for representing law-invariant risk measures using Value-at-Risk as a key component. Subsequently, we tailor our findings to derive a novel representation that adheres to the axioms of star-shapedness and cash-additivity, as expounded in Proposition 18.

Lemma 17. *Let $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ be a normalized, law-invariant risk measure. Then it admits the representation:*

$$\rho(X) = \min_{Z \in \Gamma_X} \rho \left(\sup_{\beta \in (0,1)} (VaR_\beta(X) - \gamma_Z^X(\beta)) \right).$$

Here,

$$\Gamma_X := \{Z \in \text{Dom}(\rho) : \exists Y \succeq_1 X \text{ s.t. } Y \sim \alpha Z, \alpha \in [0, 1]\},$$

and for each fixed $X \in \mathcal{X}$ and $Z \in \Gamma_X$, we define

$$\mathcal{A}_Z^X := \{\alpha \in [0, 1] : \exists Y \succeq_1 X \text{ s.t. } Y \sim \alpha Z\},$$

and the real function $\gamma_Z^X : (0, 1) \rightarrow \mathbb{R} \cup \{-\infty\}$:

$$\gamma_Z^X(\beta) := \bar{\alpha}(VaR_\beta(Z) - Z),$$

where $\bar{\alpha} = \sup \mathcal{A}_Z^X$, with that convention that if $\Gamma_X = \emptyset$ then $\gamma_Z^X = -\infty$.

Proof. We observe that γ_Z^X is well-defined for any $X \in \mathcal{X}$ and $Z \in \text{Dom}(\rho)$, as we consider $\bar{\alpha} = \sup \mathcal{A}_Z^X$, which is clearly unique. Proceeding analogously as in the proof of Proposition A.4 in [28], it can be verified that any law-invariant risk measure can be represented as:

$$\rho(X) = \min_{\substack{W \in \mathcal{X} \\ W \succeq_1 X}} \rho(W + \sup_{\beta \in (0,1)} (VaR_\beta(X) - VaR_\beta(W))).$$

We fix $X \in \text{Dom}(\rho)$, otherwise the statements are trivial. If $Z \in \Gamma_X$, then for any $\alpha \in \mathcal{A}_Z^X$ it holds that $\alpha Z \succeq_1 X$, implying that also $\bar{Y} := \bar{\alpha}Z$ verifies $\bar{\alpha}Z \succeq_1 X$. Indeed, by the properties of the supremum, there exists a sequence $(\alpha_n)_{n \in \mathbb{N}} \subseteq \mathcal{A}_Z^X$ such that $\alpha_n \rightarrow \bar{\alpha}$. Thus, for any $n \in \mathbb{N}$ it results that $Y_n \succeq_1 X$ and $Y_n \sim \alpha_n Z$. Letting $n \rightarrow \infty$, we have that $\bar{Y} := \bar{\alpha}Z$ clearly satisfies $\bar{Y} \sim \bar{\alpha}Z$ and $\bar{Y} \succeq_1 X$, so the supremum is indeed a maximum. Hence, for any $Z \in \Gamma_X$ we have

$$\rho(X) \leq \rho(\bar{\alpha}Z + \sup_{\beta \in (0,1)} (VaR_\beta(X) - \bar{\alpha}VaR_\beta(Z))). \quad (5.2)$$

Taking the minimum over $Z \in \Gamma_X$ on both members of Equation (5.2) we have:

$$\begin{aligned} \rho(X) &= \min_{\substack{W \in \mathcal{X} \\ W \succeq_1 X}} \rho(W + \sup_{\beta \in (0,1)} (VaR_\beta(X) - VaR_\beta(W))) \\ &\leq \min_{Z \in \Gamma_X} \rho(\bar{\alpha}Z + \sup_{\beta \in (0,1)} (VaR_\beta(X) - \bar{\alpha}VaR_\beta(Z))) \\ &= \min_{Z \in \Gamma_X} \rho(\sup_{\beta \in (0,1)} (VaR_\beta(X) - \gamma_Z^X(\beta))) \leq \rho(X), \end{aligned}$$

where the last inequality follows by taking $Z = X$, given that $\bar{\alpha} = 1$ attains the supremum of the set \mathcal{A}_Z^X . \square

The following proposition demonstrates that our results genuinely extend those obtained in Theorem 5 of [9]. Not only do we broaden the setting to encompass the general space \mathcal{X} , but we also establish that the minimum can be taken over a set strictly smaller than the acceptance set of ρ , denoted as $\mathcal{B}\rho := \{Z \in \mathcal{X} : \rho(Z) \leq 0\}$. More specifically, the corollary underscores that the minimum can be computed over a set $\Gamma_X \subseteq \mathcal{B}\rho$, which is contingent on the choice of $X \in \mathcal{X}$.

Proposition 18. *A functional $\rho : \mathcal{X} \rightarrow \mathbb{R} \cup \{+\infty\}$ is a normalized, law-invariant, star-shaped and cash-additive risk measure if and only if it admits the representation:*

$$\rho(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\},$$

where for each $X \in \mathcal{X}$, Γ_X is a set of random variables such that $\lambda\Gamma_X \subseteq \Gamma_{\lambda X}$ for any $\lambda \in (0, 1]$, and $\Gamma_{X+m} = \Gamma_X$ for all $m \in \mathbb{R}$. In addition, $\gamma_Z^X : (0, 1) \rightarrow \mathbb{R} \cup \{-\infty\}$ is an increasing function such that $\max_{Z \in \Gamma_0} \gamma_Z^0(0^+) = 0$. Moreover, for any $Z \in \Gamma_X$, $\lambda \in (0, 1]$ it holds that $\lambda\gamma_Z^X = \gamma_{\lambda Z}^{\lambda X}$ and for any $m \in \mathbb{R}$ it results that $\gamma_Z^{X+m} = \gamma_Z^X$. Furthermore, for each $X \in \mathcal{X}$, the set Γ_X takes the following form:

$$\Gamma_X := \{Z \in \text{Dom}(\rho) : \exists Y \succeq_1 X, Y \sim \alpha Z + c, (\alpha, c) \in [0, 1] \times \mathbb{R}, \rho(Z) \leq 0\},$$

and the function $\gamma_Z^X : (0, 1) \rightarrow \mathbb{R} \cup \{-\infty\}$ is defined by the formula:

$$\gamma_Z^X(\beta) := \bar{\alpha} VaR_\beta(Z), \quad Z \in \Gamma_X,$$

with that convention that if $\Gamma_X = \emptyset$ then $\gamma_Z^X = -\infty$. Here, $\bar{\alpha} := \sup \mathcal{A}_Z^X$ where

$$\mathcal{A}_Z^X := \{\alpha \in [0, 1] : \exists Y \succeq_1 X, c \in \mathbb{R} \text{ s.t. } Y \sim \alpha Z + c\}.$$

Proof. We start by proving the ‘only if’ implication. We assume $X \in \text{Dom}(\rho)$, otherwise the statements are trivial. For every $X \in \mathcal{X}$ and $Z \in \Gamma_X$, the function γ_Z^X is well-defined, similarly to what occurs in Lemma 17. Furthermore, we observe that $\sup \mathcal{A}_Z^X = \max \mathcal{A}_Z^X$, when $\mathcal{A}_Z^X \neq \emptyset$. It is easy to verify that if $Z \in \Gamma_X$ is non-constant and Y is non-constant, there exists a unique $(\alpha, c) \in [0, 1] \times \mathbb{R}$ such that $Y \sim \alpha Z + c$ (see the proof of Proposition 9). Thus, if \mathcal{A}_Z^X is non-empty, it consists of a single element, and the supremum is reached. When Y is constant and Z is non-constant, the only admissible choice is $(\alpha, c) = (0, Y)$, and the supremum is attained. Finally, if Z is constant, then Y must also be constant, and we can choose $(\bar{\alpha}, \bar{c}) = (1, Y - Z)$. In this case, $\bar{\alpha} = 1$ is an element of \mathcal{A}_Z^X which reaches the supremum, as $\mathcal{A}_Z^X \subseteq [0, 1]$.

Min-sup representation: First, we note that fixing $X \in \text{Dom}(\rho)$ and $Z \in \Gamma_X$ and defining

$$X \mapsto \tilde{\gamma}_Z(X) := \sup_{\beta \in (0,1)} \{(VaR_\beta(X) - VaR_\beta(\bar{\alpha}Z)) + \bar{\alpha}\rho(Z)\},$$

with $\beta \in (0, 1)$, it holds that:

$$\tilde{\gamma}_Z(X) \geq \rho(\bar{\alpha}Z + \sup_{\beta \in (0,1)} \{VaR_\beta(X) - VaR_\beta(\bar{\alpha}Z)\}).$$

Here, the inequality is due to star-shapedness and cash-additivity of ρ . We have

$$\rho(\bar{\alpha}Z + \sup_{\beta \in (0,1)} \{VaR_\beta(X) - VaR_\beta(\bar{\alpha}Z)\}) = \rho(\bar{\alpha}Z + \bar{c} + \sup_{\beta \in (0,1)} \{VaR_\beta(X) - VaR_\beta(\bar{\alpha}Z + \bar{c})\}),$$

where $\bar{c} \in \mathbb{R}$ is the element such that $Y \sim \bar{\alpha}Z + \bar{c}$. Such $\bar{c} \in \mathbb{R}$ exists since $\bar{\alpha} \in \mathcal{A}_Z^X$, as we proved in the first part of the proof. Thus, $\bar{\alpha}Z + \bar{c} \succeq_1 X$, resulting in:

$$\begin{aligned} \tilde{\gamma}_Z(X) &\geq \rho \left(\bar{\alpha}Z + \bar{c} + \sup_{\beta \in (0,1)} VaR_\beta(X) - VaR_\beta(\bar{\alpha}Z + \bar{c}) \right) \\ &\geq \min_{\substack{W \in \mathcal{X} \\ W \succeq_1 X}} \rho \left(W + \sup_{\beta \in (0,1)} (VaR_\beta(X) - VaR_\beta(W)) \right) = \rho(X), \end{aligned}$$

where the last equality follows from the proof of Lemma 17. Hence, $\min_{Z \in \Gamma_X} \tilde{\gamma}_Z(X) \geq \rho(X)$.

Since cash-additivity yields $\rho(X - \rho(X)) = 0$ it holds that $\tilde{Z} := X - \rho(X) \in \Gamma_X$ with $(\bar{\alpha}, \bar{c}) = (1, \rho(X))$. In particular, we have $\tilde{\gamma}_{\tilde{Z}}(X) = \rho(X)$, leading to:

$$\begin{aligned} \rho(X) &= \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \bar{\alpha}(VaR_\beta(Z) - \rho(Z))\} \\ &= \min_{\substack{Z \in \Gamma_X \\ \rho(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \bar{\alpha}VaR_\beta(Z)\} \\ &= \min_{\substack{Z \in \Gamma_X \\ \rho(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}. \end{aligned}$$

Furthermore, given that $Z \in \Gamma_X \implies \rho(Z) \leq 0$ it follows that:

$$\begin{aligned} \rho(X) &= \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \bar{\alpha}(VaR_\beta(Z) - \rho(Z))\} \\ &\leq \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \bar{\alpha}VaR_\beta(Z)\} \\ &\leq \min_{\substack{Z \in \Gamma_X \\ \rho(Z)=0}} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \bar{\alpha}VaR_\beta(Z)\} = \rho(X), \end{aligned}$$

hence

$$\rho(Z) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}.$$

The properties of Γ_X properties of γ_Z^X can be verified as in the proof of Corollary 16.

Now we are ready to prove the converse implication. We want to show that

$$\rho(X) = \min_{Z \in \Gamma_X} \sup_{\beta \in (0,1)} \{VaR_\beta(X) - \gamma_Z^X(\beta)\}, \quad X \in \mathcal{X},$$

is monotone, normalized, star-shaped, law-invariant and cash-additive. Normalization and star-shapedness can be established following the proof of Proposition 15, while the proof of cash-additivity closely mirrors the proof provided in Corollary 16.

Monotonicity: This property is obvious once observed that VaR_β is monotone and $\Gamma_{X_1} \subseteq \Gamma_{X_2}$ as soon as $X_1 \geq X_2$.

Law-invariance: Let us consider $X, X' \in \mathcal{X}$ such that $X \sim X'$. We need to verify that $\Gamma_X = \Gamma_{X'}$. Given that $X \sim X'$, both $X \succeq_1 X'$ and $X' \succeq_1 X$ apply, resulting in $\Gamma_X \subseteq \Gamma_{X'}$ and $\Gamma_{X'} \subseteq \Gamma_X$, which in turns lead to $\Gamma_X = \Gamma_{X'}$. This equality, along with the law-invariance of VaR_β , implies that ρ is law-invariant. \square

6 Illustrative Examples

While the prime motivation for this paper comes directly from [2, 4, 5, 30, 31], whence the aim of establishing novel characterization results for law-invariant return and star-shaped risk measures, we show in this section that such risk measures may also arise naturally from

more classical settings. Indeed, we provide three examples to illustrate the inherent star-shaped nature of certain risk measures, also when cash-additivity or cash-subadditivity are not preserved. Specifically, the examples show that during the recent extended period of negative interest rates induced by central banks' monetary policies, investors may naturally comply with non-cash-(sub)additive risk measures that remain star-shaped or even positively homogeneous.

Example 19. As shown in [19], in the absence of a zero coupon bond, the presence of ambiguity with respect to interest rates naturally leads to a relaxation of the axiom of cash-additivity, with cash-subadditivity being assumed instead. Over the last decade, the EONIA index, which tracks unsecured lending transactions in the interbank market, has assumed negative values. Consider a bank, or more generally, a financial institution, assessing the risk of an asset X_T at the present time $t = 0$, where $T > 0$ denotes the asset's time to maturity. Suppose the institution evaluates the risk of $X_T \in \mathcal{X}$ using a spot risk measure $\rho_0 : L^1 \rightarrow \mathbb{R} \cup \{+\infty\}$ defined on the discounted price $D_T X_T$, where $D_T \in \mathcal{X}^*$ represents a discount factor (for more details, see Section 2.4 in [19]). In the context of negative interest rates, the discount factor D_T can take values greater than 1. Therefore, if the interest rate is subject to ambiguity, fluctuating between two (not necessarily bounded) random variables $0 \leq D_b \leq D_u$, with $D_b, D_u \in \mathcal{X}^*$, an ambiguity-averse investor may select the risk measure as follows:

$$\tilde{\rho}(X_T) = \sup_{D_T \in \mathcal{X}^*} \{\rho_0(D_T X_T) : D_b \leq D_T \leq D_u\}.$$

Even when ρ_0 is a monetary risk measure, $\tilde{\rho}$ can be non-cash-(sub)additive, as D_T can take arbitrarily large values since D_u can be an unbounded random variable.

The fact that $D_b, D_u \in \mathcal{X}^*$ can be unbounded random variables is critical. Indeed, if the discount factors were bounded, the situation may be simplified to that of cash-subadditive risk measures. Specifically, in that case, by defining $\bar{\rho} := \frac{1}{\text{ess sup } D_u} \tilde{\rho}$, one can ensure that $\bar{\rho}$ is cash-subadditive, provided that ρ_0 is monotone and cash-(sub)additive.

A classic industry measure of risk is the Value-at-Risk. Therefore, we can consider $\rho_0 = \text{VaR}_\beta$. In this setting, the resulting risk measure is given by:

$$\tilde{\rho}(X_T) = \sup_{D_T \in \mathcal{X}^*} \{\text{VaR}_\beta(D_T X_T) : D_b \leq D_T \leq D_u\}.$$

This risk measure is neither convex nor cash-(sub)additive, but it is positively homogeneous (thus star-shaped) and law-invariant, inheriting these properties from VaR_β . In particular, the thesis of Proposition 15 concerning positively homogeneous functionals can be applied to represent $\tilde{\rho}$ as in Equation (5.1).

Example 20. In the same context as in Example 19, we consider a generalization of VaR_β that is closely connected to the concept of Λ -VaR, as shown in Theorem 3.1 of [28]. For a thorough discussion of Λ -VaR risk measures, we refer to [23]. Let us fix $x \in \mathbb{R}$ and define:

$$\begin{aligned} \tilde{\rho}(X_T) &= \sup_{D_T \in \mathcal{X}^*} \{\text{VaR}_\beta(D_T X_T) \vee x : D_b \leq D_T \leq D_u\} \\ &= \sup_{D_T \in \mathcal{X}^*} \{\text{VaR}_\beta(D_T X_T) : D_b \leq D_T \leq D_u\} \vee x. \end{aligned}$$

Since the maximum operation preserves star-shapedness, and both VaR_β and the trivial risk measure $\rho(X) = x$ for any $X \in \mathcal{X}$ are star-shaped, the resulting risk measure $\tilde{\rho}$ is law-invariant and (genuinely) star-shaped. However, in this case, positive homogeneity does not hold in general. Once again, $\tilde{\rho}$ is neither convex nor cash-(sub)additive. The parameter x can be interpreted as follows: it represents a barrier below which the risk evaluation corresponding to the asset X_T cannot fall, possibly due to market frictions or other constraints. Consequently, the investor is required to retain the larger of two amounts — the assessment of the risk linked to the asset X_T as determined by the Value-at-Risk and the minimum threshold $x \in \mathbb{R}$. The fixed amount x remains constant and is unrelated to the value of X_T ,

but instead may be influenced by external factors. In this case, it is also possible to derive a representation of $\tilde{\rho}$ akin to the one presented in Equation (5.1). It should be noted that $\tilde{\rho}$ is not normalized. However, the representation results remain valid, as stated at the outset of Section 5.

Example 21. Consider a firm that wishes to establish an insurance contract to cover the risk associated with an asset X_T . Let A represent the set of insurance companies accessible to the firm. Suppose each insurer has its own spot monetary, convex, and normalized risk measure ρ_0^a with $a \in A$ to assess the risk of X_T . Furthermore, suppose the cost of each contract is equal to the risk assessment made by the insurer through its risk measure. Thus, for each asset X_T , the firm will choose to pay the minimum amount:

$$\rho(X_T) := \min_{a \in A} \sup_{D_T \in \mathcal{X}^*} \{\rho_0^a(D_T X_T) : D_b^a \leq D_T \leq D_u^a\}.$$

Here, each insurer has its own perspective on ambiguous interest rates, allowing D_T to vary between D_b^a and D_u^a , with $D_b^a, D_u^a \in \mathcal{X}^*$ such that $0 \leq D_b^a \leq D_u^a$, and where $a \in A$. The resulting risk measure is star-shaped but not convex in general. Furthermore, as observed in the examples above, ρ lacks cash-(sub)additivity. If the risk measures employed by the insurance companies are SSD-consistent, such as in the case of Expected Shortfall, the resulting risk measure ρ inherits this property. Under this circumstance, if \mathcal{X} is a rearrangement-invariant space, we can represent ρ as in Equation (4.1).

These three examples underscore the significance of establishing a comprehensive framework for law-invariant (or SSD-consistent) star-shaped risk measures, which remains applicable also when cash-(sub)additivity is not preserved.

References

- [1] ACCIAIO, B. (2007). Optimal risk sharing with non-monotone monetary functionals. *Finance and Stochastics* 11, 267-289.
- [2] AYGÜN, M., F. BELLINI AND R. J. A. LAEVEN (2023). Elicitability of return risk measures. *Preprint on arXiv:2302.13070v2*.
- [3] BÄUERLE, N. AND A. MÜLLER (2006). Stochastic orders and risk measures: Consistency and bounds. *Insurance: Mathematics and Economics* 38, 132-148.
- [4] BELLINI F., P. KOCH-MEDINA, C. MUNARI AND G. SVINDLAND (2021). Law-invariant functionals on general spaces of random variables. *SIAM Journal on Financial Mathematics* 12, 318-341.
- [5] BELLINI, F., R. J. A. LAEVEN AND E. ROSAZZA GIANIN (2018). Robust return risk measures. *Mathematics and Financial Economics* 12, 5-32.
- [6] BELLINI F., R. J. A. LAEVEN AND E. ROSAZZA GIANIN (2021). Dynamic robust Orlicz premia and Haezendonk-Goovaerts risk measures. *European Journal of Operational Research* 12, 5-32.
- [7] BIAGINI, S. AND M. FRITTELLI (2008). A unified framework for utility maximization problems: An Orlicz space approach. *The Annals of Applied Probability* 18, 929-966.
- [8] BIAGINI, S. AND M. FRITTELLI (2009). On the extension of the Namioka-Klee Theorem and on the Fatou property for risk measures. *Optimality and Risk—Modern Trends in Mathematical Finance*, 11-28.
- [9] CASTAGNOLI, E., G. CATTELAN, F. MACCHERONI, C. TEBALDI AND R. WANG (2022). Star-shaped risk measures. *Operations Research* 70, 2637-2654.

- [10] CERREIA-VIOGLIO, S., F. MACCHERONI, M. MARINACCI AND L. MONTRUCCHIO (2011). Risk measures: Rationality and diversification. *Mathematical Finance* 21, 743-774.
- [11] CHERIDITO, P. AND T. LI (2009). Risk measures on Orlicz hearts. *Mathematical Finance* 19, 189-214.
- [12] CHONG, K. M. AND N. M. RICE (1971). Equimeasurable rearrangements of functions. *Queen's Papers in Pure and Applied Mathematics* 28, 1-177.
- [13] DANA, R. A. (2005). A representation result for concave Schur concave functions. *Mathematical Finance* 15, 613-634.
- [14] DELBAEN, F. (2002). Coherent risk measures on general probability spaces. *Essays in Honour of Dieter Sondermann*, Springer, Berlin, Heidelberg.
- [15] DELBAEN, F. (2006). The structure of m -stable sets and in particular of the set of risk neutral measures. In: Émery, M. and M. Yor (eds.). *In Memoriam Paul-André Meyer, Lecture Notes in Mathematics* 1874, 215-258, Springer, Berlin, Heidelberg.
- [16] DELBAEN, F. (2012). *Monetary Utility Functions*. Osaka University Press, Osaka.
- [17] DEPREZ, O. AND H. U. GERBER (1985). On convex principles of premium calculation. *Insurance: Mathematics and Economics* 4(3), 179-189.
- [18] DRAPEAU, S. AND M. KUPPER (2013). Risk preferences and their robust representation. *Mathematics of Operations Research* 38, 28-62.
- [19] EL KAROUI, N. AND C. RAVANELLI (2009). Cash sub-additive risk measures and interest rate ambiguity. *Mathematical Finance* 19, 561-590.
- [20] FILIPOVIC, D. AND G. SVINDLAND (2008). Optimal capital and risk allocations for law- and cash-invariant convex functions. *Finance and Stochastics* 12, 423-439.
- [21] FÖLLMER, H. AND A. SCHIED (2002). Convex measures of risk and trading constraints. *Finance and Stochastics* 6, 429-447.
- [22] FÖLLMER, H. AND A. SCHIED (2011). *Stochastic Finance*. 3rd ed., De Gruyter, Berlin.
- [23] FRITTELLI, M., M. MAGGIS AND I. PERI (2013). Risk measures on $\mathcal{P}(\mathbb{R})$ and value at risk with probability/loss function. *Mathematical Finance* 24, 442-463.
- [24] FRITTELLI, M. AND E. ROSAZZA GIANIN (2002). Putting order in risk measures. *Journal of Banking and Finance* 26, 1473-1486.
- [25] FRITTELLI, M. AND E. ROSAZZA GIANIN (2005). Law-invariant convex risk measures. *Advances in Mathematical Economics* (pp. 33-46). Springer, Tokyo.
- [26] GOOVAERTS, M. J., F. E. C. DE VYLDER AND J. HAEZENDONCK (1984). *Insurance Premiums*. North-Holland Publishing, Amsterdam.
- [27] HAEZENDONCK, J. AND M. J. GOOVAERTS (1982). A new premium calculation principle based on Orlicz norms. *Insurance: Mathematics and Economics* 1, 41-53.
- [28] HAN X., Q. WANG, R. WANG AND J. XIA (2022). Cash-subadditive risk measures without quasi-convexity. *Preprint on arXiv:2110.12198v4*.
- [29] KUSUOKA, S. (2001). On law-invariant coherent risk measures. *Advances in Mathematical Economics* (pp. 83-95). Springer, Tokyo.
- [30] LAEVEN, R. J. A. AND E. ROSAZZA GIANIN (2022). Quasi-logconvex measures of risk. *Preprint on arXiv:2208.07694v1*.

- [31] LAEVEN, R. J. A., E. ROSAZZA GIANIN AND M. ZULLINO (2023). Dynamic return and star-shaped risk measures via BSDEs. *Preprint on arXiv:2307.03447v2*.
- [32] LAEVEN, R. J. A. AND M. A. STADJE (2013). Entropy coherent and entropy convex measures of risk. *Mathematics of Operations Research* 38, 265-293.
- [33] LAEVEN, R. J. A. AND M. A. STADJE (2014). Robust portfolio choice and indifference valuation. *Mathematics of Operations Research* 39, 1109-1141.
- [34] MACHINA, M. J. AND D. SCHMEIDLER (1992). A more robust definition of subjective probability. *Econometrica* 60, 745-780.
- [35] MAO, T. AND R. WANG (2020). Risk aversion in regulatory capital calculation. *SIAM Journal on Financial Mathematics* 11(1), 169-200.
- [36] MARINACCI, M. (2002). Probabilistic sophistication and multiple priors. *Econometrica* 70, 755-764.
- [37] MCNEIL, A. J., R. FREY AND P. EMBRECHTS (2015). *Quantitative Risk Management: Concepts, Techniques and Tools*. Revised Edition. Princeton, NJ: Princeton University Press.
- [38] PFLUG, G. CH., AND W. RÖMISCH (2007). *Modeling, Measuring and Managing Risk*. World Scientific, Singapore.
- [39] SHAKED, M. AND J. G. SHANTHIKUMAR (2007). *Stochastic Orders*. Springer Series in Statistics.
- [40] SVINDLAND, G. (2008). *Convex Risk Measures Beyond Bounded Risk*. Ph.D. Thesis, Ludwig Maximilian Universität München.
- [41] WANG, R., Y. WEI AND G. E. WILLMOT (2020). Characterization, robustness and aggregation of signed Choquet integrals. *Mathematics of Operations Research* 45(3), 993-1015.